

GUANYLYL CYCLASES ACROSS THE TREE OF LIFE

Pauline Schaap

School of Life Sciences, University of Dundee, UK

TABLE OF CONTENTS

1. Abstract
2. Introduction
3. Animals
 - 3.1. The cyclase catalytic domain
 - 3.2. Vertebrate cGMP signaling, a brief survey
 - 3.3. Invertebrate guanylyl cyclases
4. Fungi
5. Dictyostelids
6. Plants
 - 6.1. Flowering plants
 - 6.2. Chlorophyte green algae
7. Alveolates
8. Discicristates
9. Prokaryotes
10. Summary and perspective
11. Acknowledgement
12. References

1. ABSTRACT

This review explores the origins, diversity and functions of guanylyl cyclases in cellular organisms. In eukaryotes both cGMP and cAMP are produced by the conserved class III cyclase domains, while prokaryotes use five more unrelated catalysts for cyclic nucleotide synthesis. The class III domain is found embedded in proteins with a large variety of membrane topologies and other functional domains, but the vertebrate guanylyl cyclases take only two forms, the receptor guanylyl cyclases with single transmembrane domain and the soluble enzymes with heme binding domain. The invertebrates additionally show a soluble guanylyl cyclase that cannot bind heme, while the more basal metazoans may lack the heme binding enzymes altogether. Fungi, the closest relatives of the metazoans, completely lack guanylyl cyclases, but they appear again in the Dictyostelids, the next relative in line. Remarkably, the two Dictyostelid guanylyl cyclases have little in common with the vertebrate enzymes. There is a soluble guanylyl cyclase, which shows greatest sequence and structural similarity to the vertebrate soluble adenylyl cyclase, and a membrane-bound form with the same configuration as the dodecahedral adenylyl cyclases of vertebrates. There is a difference, the pseudosymmetric C1 and C2 catalytic domains have swapped position in the *Dictyostelium* enzyme. Unlike the vertebrate guanylyl cyclases, the *Dictyostelium* enzymes are activated by heterotrimeric G-proteins. Swapped C1 and C2 domains are also found in the structurally similar guanylyl cyclases of ciliates and apicomplexans, but these enzymes additionally harbour an amino-terminal ATPase module with ten transmembrane domains. G-protein regulation could not be demonstrated for these enzymes. Higher plants lack class III cyclase domains, but an unexplored wealth of guanylyl cyclases is present in the

green alga *Chlamydomonas*. Progenitors of all structural variants of the eukaryote guanylyl cyclases are found among the prokaryote adenylyl cyclases. This and the close similarity of many guanylyl cyclases to adenylyl cyclases suggests a paraphyletic origin for the eukaryote enzymes with multiple events of conversion of substrate specificity.

2. INTRODUCTION

Stimulus-induced production of the second messenger cGMP controls a broad spectrum of physiological responses in vertebrates, such as vision, olfaction, smooth muscle relaxation and homeostasis of electrolyte and fluid levels in blood and intestine. The cGMP-mediated functions show only limited overlap with those mediated by the chemically related messenger cAMP. Metazoans have developed sets of proteins with different membrane topologies and domain architectures for synthesis of either cAMP or cGMP. As a consequence, the activation mechanisms of guanylyl cyclases bear little in common with those of adenylyl cyclases. The strict separation into architecturally different forms is quite remarkable in view of the fact that the catalytic domains of adenylyl- and guanylyl cyclases are very similar and can be interconverted by substitution of only a few amino-acids.

cGMP has also been implicated in controlling motility in protists, such as the Alveolates and Dictyostelids. These organisms show a novel repertoire of architectural designs for guanylyl cyclases, accompanied by different modes of regulation. The ongoing and finished sequencing projects of many protist genomes are also uncovering new guanylyl cyclase forms.

Guanylyl cyclase diversity

ACVC1	473	ILFADIEGFTSLASQC----	TAQELVMTLNELFARFDKLAA---	ENHCLRIKILGDCYYC	525
		h M h	h h	h M h	
ACIIC2	887	VMFASIPDFKEFYTESDVNKEGLECLRL	NEIIADFDDLLSKPKFSGVEKIKTIGSTYMA	946	
DdACG	397	VFFLDIAGFTTRFSSIH----	SPEQVIQVLIKIFNSMDLLCA---	KHGIEKIKTIGDAYMA	449
RetGC	881	LYFSDIVGFTTISAMS----	EPIEVVDLLNDLYTLFDAIIG---	SHDVYKVETIGDAYMV	933
777.3	337	IMFTDIVGFTNMCHEC----	QPLEVMAFLNDLYSRFDGLVD---	IYKVYKVETIGDCYMV	389
				A/G F	
ACVC1	561	NVNMRRVGIHSGRVHCGVL-GLR	KWQFDVWSNDVTLANHMEAGGKAGRIHITKATLSYLN	619	
		P	F h M		
ACIIC2	993	DFKLRVGINHGPVIAGVI-GA	QKPQYDIWGNVTNVASRMDSTGVLDKIQVTEETSLILQT	1051	
			A/Ghh T T		
DdACG	487	GLQVRVGIHCGPVISGVISGYAKPHFD	VWGDVTNVASRMESTGIAGQIHVS	DRVYQLGKE	546
RetGC	972	PVRIRIGLHSGPCVAGVV-GLT	MPRYCLFGDVTNTASRMESTGLPYRIHVN	LSTVGILRA	1030
777.3	439	PVRLRVGLHSGPVTSGVV-GDR	MPRFCLFGDVTNVASRMESTCRPGHIHVS	AATQARLPN	487
AtGC1	20	-----G-ILDGN-GDSTFPK	YCLFDDPLVSDGKYRDAGLPSSSHMD-----	58	

Figure 1. Alignment of cyclase catalytic domains. The catalytic domains of the mammalian ACV C1 domain and ACII C2 domains, mammalian RetGC, *Dictyostelium discoideum* ACG, *Chlamydomonas reinhardtii* gene 777.3 and *Arabidopsis thaliana* AtGC1 were aligned using Clustal W (146). Residues essential for substrate binding and catalysis are highlighted. Green: residues that form hydrophobic pocket; red/yellow: residues that determine guanine/adenine specificity; blue: residues that coordinate the Mg^{2+} ions; amber: interaction with forskolin; magenta: residues that stabilize the transition state; tan: residues that interact with Ppi. Genbank accession: CfACV: AAC32726; RrACII A41541; DdACG: Q03101; HsRetGC: NP_000171; AtGC1: AAM51559. The gene.777.3 sequence can be found at <http://genome.jgi-psf.org/chlre1/chlre1.home.html>

This review summarizes the evidence for the existence of cGMP signaling in the eight major group of eukaryotes and in the prokaryotes and compares the structure and function of their guanylyl cyclases.

3. ANIMALS

3.1. The cyclase catalytic domain

In all characterized eukaryote guanylyl- and adenylyl cyclases, cyclic nucleotide synthesis is carried out by the conserved class III cyclase domain. The crystal structure of the catalytic core of the G-protein stimulated vertebrate adenylyl cyclase is resolved and this also provides insight in the catalytic mechanism of the guanylyl cyclases. G-protein activated adenylyl cyclases harbour two sets of six transmembrane helices that are separated by the cyclase C1 domain and followed by the cyclase C2 domain. These domains form an antiparallel dimer with ATP binding to the dimer interface. Potentially, two ATP molecules, each associated with two Mg^{2+} ions can be bound. However, the C1-C2 dimer accommodates only one ATP, because the C2 domain lacks two aspartate residues that should coordinate the Mg^{2+} ions, while C1 lacks an arginine and asparagine pair that interacts with the phosphoryl moiety of the same ATP and stabilizes the transition state (1-4). The adenine ring interacts with a conserved lysine, a glutamine and an aspartate residue. Guanylyl cyclases carry a glutamate, an arginine and a cysteine at the equivalent positions (figure 1). Mutagenesis of these residues into their adenylyl cyclase counterpart alters the substrate preference of both soluble- and receptor guanylyl cyclase from GTP to ATP (5, 6). For those adenylyl- and guanylyl cyclases that harbour a single catalytic domain and form homodimers, all essential residues have to be present in the single domain. In these enzymes two symmetric ATP or GTP binding sites are created (7, 8). The C1-C2 dimer accommodates the diterpene forskolin at the position of the

degenerate ATP binding site. Forskolin is a well-known activator of mammalian adenylyl cyclases, and it does so by stabilizing the association of the C1 and C2 monomers. It interacts with several hydrophobic residues in a deep hydrophobic cleft, and with a serine residue that forms a hydrogen bond with the 7-acetyl group of forskolin. This serine replaces one of the Mg^{2+} binding aspartates in C2 (1, 2). $G_{s\alpha}$, the physiological activator, also binds simultaneously to the C1 and C2 monomers and activates catalysis by stabilizing the catalytically competent juxtaposition of the monomers (1).

3.2. Vertebrate cGMP signaling, a brief survey

There are two topologically distinct guanylyl cyclases in mammalian cells, the receptor guanylyl cyclases and the soluble guanylyl cyclases, neither of which is activated by G-proteins. The receptor guanylyl cyclases are integral transmembrane proteins with a single transmembrane helix that separates the external ligand binding domain from a carboxyl-terminal region that contains a protein kinase homology domain (KHD), a coiled-coil (CC) domain and a single guanylyl cyclase (GC) domain (figure 2). The majority of these proteins already form dimers in the inactive state by interactions between their CC and KHD domains (9-11). Binding of the ligand causes juxtamembrane closure of the extracellular domain, which reorients the intracellular KHD domain. This supposedly allows it to bind ATP, which in turn triggers a second conformational change, which results in activation of the cyclase domain (12-14). The ligands are usually peptides, such as the natriuretic peptides for GC-A and GC-B (10, 15, 16) and enterotoxins and guanylin for GC-C (17-19), or unknown odorants for GC-D (20).

The Ca^{2+} regulated retinal guanylyl cyclases (RetGCs) are not (known to be) regulated by ligand; their dimerization state and therefore their activity depends on




Figure 2. Domain architectures of guanylyl- and adenylyl cyclases. The amino-acid sequences of the indicated proteins were analysed with SMART (147) for SMART and PFAM functional domains (148) and by TMHMM (149) for putative transmembrane domains. The domain architecture of all proteins printed in bold is presented at the correct relative scale. The proteins indicated between brackets contain a similar array of domains, but the relative scale may be different. Abbreviations: ANP-R: atrial natriuretic peptide receptor; KHD: kinase homology domain; AAA: ATPase family associated with various cellular activities (76); PP2C: Serine/threonine phosphatases, family 2C; RA: Ras association domain; CHASE: cyclase and histidine kinase associated extracellular sensor; HK: histidine kinase; HA: histidine kinase like ATP-ase; RR: response regulator; C39: peptidase C39 family; VGC: voltage-gated channel; BLUF: sensors of blue-light using FAD; GAF: domain present in phytochromes and cGMP phosphodiesterases. Genbank accession numbers: HsGCA: AAH63304; HsRetGC: NP_000171; S.purGC: P16065; HssolGC α : NP_000847; HssolGC β : NP_000848; MsGC-I: AAC62238; HsAC1: NP_066939; HssAC1: NP_060887; ScCYR1: P08678; DdsGC: AAK92097; DdGCA: CAB42641; DdACA: AAA33163; DdACG: Q03101; DdAcrA: AAD50121; AtGC1: AAM51559; ZmPsiP: CAC59976; P.tetGC: PTE238859; P.falGC α : PFA245435; P.tetAC: CAD60412; P.falAC1: AAO64441; P.falAC2: NP_704518; T.bruESAG4: Q26721; T.cruADC1: CAA09919; L.donRacA: Q27675; EgAC α : BAB85619; Syn.spp Cya2: NP_440289; Nostoc CyaC: NP_489003.

interaction of each monomer with a Guanylyl Cyclase Activating Protein (GCAP), which dimerizes in the absence, but not in the presence of Ca^{2+} (21-23). The RetGCs nevertheless harbour the conserved ligand binding, KHD and CC domains of the other receptor guanylyl cyclases (figure 2).

The soluble guanylyl cyclases (SolGCs) form a heterodimer of two isoforms, α and β , which each carry a guanylyl cyclase domain at the C-terminus, and a region at the N-terminus that coordinates a single heme group per dimer (24-26). Two cysteine residues (Cys^{78} and Cys^{214}) and a histidine (His^{105}) on the β -isoform are particularly important for binding to and coordination of the heme group (27, 28). Similar to the C1 and C2 adenylyl cyclase domains, the α - and β - guanylyl cyclase domains are not identical. Like C2, the β -isoform lacks residues to coordinate the Mg^{2+} ions, while the α -isoform lacks residues that stabilize the transition state (figure 1). Therefore, also here only one catalytic site can be formed (7). SolGCs are 200-fold activated by the interaction of NO radicals with heme Fe^{2+} (29, 30). This activation is blocked by a specific inhibitor, 1H-[1,2,4]oxadiazolo[4,3-a]-quinoxalin-1-one (ODQ), which competes with NO for binding to heme and oxidizes the Fe^{2+} (31, 32). SolGCs play decisive roles in smooth muscle relaxation and are also implicated in synaptic transmission and blood platelet aggregation (26).

cGMP is detected in vertebrates by either cGMP-dependent protein kinases (PKGs) or by cGMP-gated ion channels (33, 34). PKG harbours two intramolecular cGMP binding sites, that are very similar in sequence and structure to the cAMP binding sites on the regulatory subunit of PKA (35). One copy of this canonical cNMP binding domain is present on the cGMP- (and cAMP) gated ion channels. cGMP can also bind to the GAF domain, a family of small molecule binding domains with multiple functions. The cGMP binding GAF domains are present on a number of vertebrate cyclic nucleotide phosphodiesterases, where they mediate allosteric activation of the catalytic activity by cGMP (36). There are 11 families of vertebrate phosphodiesterases, which all share the same highly conserved HD domain for cyclic nucleotide hydrolysis (37). However, the families differ in substrate selectivity, inhibitor sensitivity and the presence of GAF- and other functional domains (38, 39).

3.3. Invertebrate guanylyl cyclases

The receptor guanylyl cyclases were first found in the echinodermata (sea urchins), where they are expressed in sperm cells to detect chemotactic peptides secreted by eggs (40). More recently, they have also been identified in more basal phyla. They are very abundant in the nematode, *C.elegans*, where they are involved in olfaction (41-43) and in *Drosophila* where their function has not yet been resolved (44). A Ca^{2+} inhibited RetGC type receptor guanylyl cyclase was identified in another arthropod, *Manduca sexta* (45). The NO-stimulated soluble guanylyl cyclases are also functionally conserved in the arthropods (46, 47).

In addition to these “classical” guanylyl cyclases,

the invertebrates present us with a few new configurations. The *M.sexta* enzyme MsGC-I, shares greatest sequence similarity with the mammalian enzyme receptor guanylyl cyclase GC-B in its cyclase and coiled-coil domain, but lacks the extracellular-, transmembrane- and kinase homology domain of GC-B (figure 2). When expressed in COS-7 cells, it is soluble and active as a homodimer (48). Another *Manduca* enzyme MsGC- β 3 is closely related to the SolGC β -isoform. However, it lacks the two cysteine residues that are essential for heme binding (28) and it is active as an NO insensitive homodimer even in the presence of the MsGC- α isoform (49, 50). All predicted *C.elegans* SolGC proteins also lack the essential cysteine residues for heme binding and therefore stimulation by NO (43, 51). In addition, the gene for NO synthase is absent from the *C.elegans* genome (52). Since the Pseudocoelomate Nematoda represents a more basal metazoan form than the Eucelomate Arthropoda, this suggests that NO regulation of guanylyl cyclase activity is an evolutionary novelty of the more derived metazoans. However, this suggestion requires support from fully sequenced genomes of more basal metazoan taxa.

4. FUNGI

The fungi are a large group of non-motile organisms, which feed by adsorption of nutrients. They exist primarily as filamentous networks, hyphae, that infiltrate their substrata. The fungi are the closest relatives of the animals and some unicellular allies, with whom they form the major group of ophisthokonts (figure 3). All fungi share a highly conserved cAMP signaling pathway, that controls differentiation and pathogenicity in response to nutrient status (53, 54). The single fungal adenylyl cyclase, CYR1, is a soluble protein, that is associated to the plasma membrane by a lipid anchor (55). In addition to a single cyclase domain, CYR1 harbours a phosphatase 2C domain, a Ras Association domain and a series of leucine-rich repeats (figure 2). The latter two regions are required for binding to the monomeric G-protein Ras, which activates the enzyme. CYR1 is also activated by Gpa2, the α -subunit of a heterotrimeric G-protein (56-58). Despite, the latter resemblance to regulation of vertebrate adenylyl cyclases, the fungal cyclase domain bears greater sequence similarity to that of the *Trypanosomid* receptor adenylyl cyclases than to the vertebrate enzymes (59).

cAMP is detected by a single PKA regulatory subunit, which can interact with up to three catalytic subunits (60, 61). There are two cAMP phosphodiesterases; Pde1 is a low affinity enzyme, which harbours the HxHxDH metal binding motif of the protist type II phosphodiesterases (62), while Pde2 is a high affinity enzyme, with the HD catalytic motif of the vertebrate type I PDEs (37, 63). No genes encoding either guanylyl cyclases, cGMP binding proteins or cGMP-PDEs have been found in the completed genomes of *Saccharomyces cerevisiae* (64) and *Candida albicans* (65), nor have they been reported in other fungi. This, and the paucity of fungal adenylyl cyclases contrasts strongly with the abundance of guanylyl- and adenylyl cyclases in the related metazoans and dictyostelids.




Figure 3. Consensus phylogeny of eukaryotes. Based on molecular and ultrastructural data, most of the presently known eukaryotes can be assigned to one of eight major groups (150, 151). Courtesy of Dr. Sandra L. Baldauf.

5. DICTYOSTELIDS

The Dictyostelids are the most thoroughly studied representatives of the amoebazoa, the closest sister group of the opisthokonts (figure 3). Most members of the amoebazoa are unicellular predators that feed on bacteria, although some, like the Entamoebas, have adopted an endoparasitic life style. In addition to a unicellular feeding stage, the Dictyostelids aggregate to form a multicellular structure in response to starvation. During this process, extracellular cAMP acts as a chemoattractant to bring starving cells together. cAMP also triggers a transient increase in cytosolic cGMP (66), and mutants with defective cGMP metabolism or detection have implicated cGMP as a regulator of the chemotactic response (67-69). The *Dictyostelium* adenylyl- and guanylyl cyclases show a marked similarity to the vertebrate cyclases, but structurally similar proteins often have opposite substrate specificities (figure 2). There is both an adenylyl cyclase, ACA, and a guanylyl cyclase, GCA with the same dodecahedral topology as the G-protein regulated mammalian adenylyl cyclases (70, 71). In case of GCA, the order of the C1 and C2 domains is reversed, with C2 N-terminal to C1.

The second *Dictyostelium* guanylyl cyclase, sGC (72), is both with regard to domain architecture and sequence similarity the closest homologue of the

mammalian soluble adenylyl cyclase sAC (73, 74). Both enzymes lack putative transmembrane helices and share highly conserved C1 and C2 domains and an AAA-domain with P-loop motif (75, 76). Quite strikingly, there is no sAC in invertebrates, such as *C.elegans* and *Drosophila* (72).

In the eukaryotes, the sAC and sGC cyclase domains are most similar to those of the *D.discoideum* adenylyl cyclase, AcrA and the alveolate (*Plasmodium*, *Paramecium* and *Tetrahymena*) adenylyl cyclases. However, these enzymes are all transmembrane proteins with a single cyclase domain, and they also lack the AAA domain. As a group, sAC, sGC, AcrA and the Alveolate enzymes show much greater similarity to bacterial adenylyl cyclases, than to any other eukaryote enzyme (72, 73, 77-80).

In contrast to sAC and to all vertebrate guanylyl cyclases, sGC and GCA require a heterotrimeric G-protein, G2, for activity. However, for both proteins, the activation by G2 is indirect and may additionally involve monomeric G-proteins (81, 82). All sGC activity appears to be cytosolic, when measured in the presence of Mn^{2+} ions. However, with Mg^{2+} and the G-protein activator, GTP γ S, a much lower activity is exclusively detectable in the membrane fraction, suggesting that the G-protein regulated

population of sGC molecules is membrane-associated (72).

Dictyostelium cells express a third adenylyl cyclase, ACG, that displays the same topology as the receptor guanylyl cyclases with a large extracellular region, a single transmembrane domain and a cytosolic cyclase domain (15, 70). However, there is no coiled-coil or kinase homology domain in ACG. Furthermore, its extracellular region harbours a CHASE domain instead of the metazoan ANP-R domain (figure 2). The CHASE domain is a predicted sensor domain, that is found on some histidine kinases and bacterial adenylyl cyclases (83). ACG is active as a homodimer and harbours an intramolecular osmosensor that mediates activation of catalytic activity by high osmolarity (84). It is not yet clear whether this osmosensor is identical with the CHASE domain.

The *Dictyostelium* genome has not yielded a mammalian type PKG, but four proteins with PKG-like cNMP binding domains were identified. Two of these proteins, PdeD and PdeE harbour two cNMP binding domains each, and a metallo- β -lactamase domain. This domain is found in a number of Zn²⁺-dependent hydrolases (85) and though otherwise dissimilar from the type II cAMP-PDEs of yeast and *Dictyostelium* (62, 86), it does carry their conserved HxHxDH motif. The metallo- β -lactamase domain of PdeD encodes a highly specific cGMP-phosphodiesterase activity and that of PdeE a less specific cAMP-PDE activity. The cNMP binding domains of PdeD mediate allosteric activation of the cGMP-PDE activity by cGMP, while PdeE is allosterically activated by cAMP (87-89). This differs markedly from the vertebrate phosphodiesterases, where the GAF domain mediates allosteric activation of hydrolytic activity (36).

The other two proteins GbpC and GbpD are also multidomain proteins and most likely the sole effectors for cGMP (90). Both contain two cGMP binding domains, a GRAM domain, a RasGEF and a RasGEF-associated domain. GbpC additionally harbours a MAPKKK domain, a Ras domain and a DEP domain. The only known proteins with cNMP binding domains, that share at least some of this plethora of domains are the vertebrate EPACs, in which a RapGEF domain is activated by the associated cAMP binding domain (91). GbpC and GbpD are considered to regulate the phosphorylation status of myosin II in the chemotactic response (69).

6. PLANTS

6.1. Flowering plants

The existence of cyclic nucleotide signaling in higher plants has been controversial for many years. Compared to organisms with well-defined functions for cyclic nucleotides, the levels of both cAMP and cGMP in higher plants are very low and it has been challenging to convincingly demonstrate the presence of nucleotidyl cyclase and phosphodiesterase activities (92). The completed genome sequence of the *Arabidopsis thaliana* does not contain sequences for the universal class III nucleotidyl cyclases (93). However, two unusual cyclase genes were identified. The *Zea mays* gene *PSiP* was

identified as a putative adenylyl cyclase on the basis of the homology of its leucine-rich repeat region to that of the fungal ACs (94). *PSiP* also harbours an AAA type ATP-ase domain, but there is no class III cyclase domain. Its function as an adenylyl cyclase is indicated by the fact that *PSiP* can complement defective sugar fermentation and cAMP production in an *E.coli* adenylyl cyclase null mutant. cAMP production is stimulated by forskolin, which also stimulates the process of pollen tube outgrowth that is dependent on *PSiP* (94). In view of the fact that forskolin interacts in a highly structure-specific manner at the interface of the C1 and C2 domains of not even all vertebrate cyclases (1, 2), this stimulation of *PSiP*, which does not harbour a class III cyclase domain is rather remarkable.

A motif search of the *Arabidopsis* genome with conserved amino-acids in the guanylyl cyclase catalytic centre yielded a candidate protein, AtGC1, that raises cGMP levels when expressed in *E.coli* and displays Mg²⁺ dependent GC activity when purified (95). AtGC1 can also produce cAMP at a three-fold lower rate. Only the posterior third of the GC catalytic domain is conserved at the N-terminus of AtGC1, which means that several residues for GTP and Me²⁺ binding are lacking (figures 1, 2). However, additional functionality may reside in the C-terminal region of the protein. AtGC1 is a soluble protein that harbours none of the heme- or ligand binding, coiled-coil or kinase homology domains that are common to vertebrate guanylyl cyclases (15, 95, 26).

Quite fascinatingly, plants secrete extracellular peptides, irPNP's, that bind to antibodies against mammalian atrial natriuretic peptide (ANP) and share a small region of high sequence identity with ANPs (96). Both ANP and irPNP regulate plant solute homeostasis by inducing opening of leaf stomatal pores and ion channels in roots (97-99). These effects can be mimicked by 8Br-cGMP and are in case of the stomatal pores accompanied by an increase in cGMP levels (100).

NO, the activator of the vertebrate SolGC, also regulates important processes in plant physiology and development (101) and increases cGMP levels in spruce and tobacco (102, 103). Inhibitors of vertebrate SolGC, such as ODQ and LY83583 antagonize some NO-induced responses and 8Br-cGMP can reverse this inhibition (103-105) and in one case also mimick the effect of NO (103). If AtGC1 is the only plant guanylyl cyclase, it has to be assumed that both the effects of NO and the natriuretic peptides are mediated by auxiliary sensor proteins that interact with AtGC1. This would represent a truly remarkable case of convergent evolution.

No genes encoding either PKG or the regulatory subunit of PKA have been found in the *Arabidopsis* genome, but the characteristic cyclic nucleotide (cNMP) binding domain of these proteins is present in a large family of plant ion-channels. These channels show a similar membrane topology as the vertebrate cyclic nucleotide gated ion channels; only the localization of the additional calmodulin binding domain is different. In the

vertebrate channels this domain is located at the N-terminus, while in the plant channels it overlaps with the cNMP binding domain at the C-terminus (106-110). Most, if not all, plant cNMP domains harbour the Ala to Ser/Thr substitution that marks cGMP specificity (35, 107, 111).

Only a few members of the plant cyclic nucleotide gated channel (CNGC) family have been investigated. AtCNGC2/DND1 mediates pathogen-induced apoptosis and displays inwardly rectified transport of K^+ , Ca^{2+} and other cations except Na^+ . Channel opening is activated equally well by 8Br-cAMP and 8Br-cGMP (112-115). AtCNGC4/HLM1 is also involved in defence to pathogens, but is permeable to both K^+ and Na^+ . Here, cGMP triggers channel opening more efficiently than cAMP (116). The presence of cyclic nucleotide gated ion-channels is thus far the strongest genetic evidence that flowering plants use cyclic nucleotides and perhaps particularly cGMP as signaling molecules.

6.2. Chlorophyte green algae

The chlorophyte green algae are basal in the line of descent that gave rise to the flowering plants (figure 3) and the unicellular alga *Chlamydomonas* is used as a model system for plant signaling and cell biology. cAMP signaling plays a well-documented role in the mating process of *Chlamydomonas*, where it is produced when cells of opposite mating type agglutinate. cAMP is produced by an adenylyl cyclase activity, that is insensitive to regulators of vertebrate adenylyl cyclases, such as forskolin and GTP γ S. The cells also display a cAMP phosphodiesterase activity, that is inhibited by IBMX, an inhibitor of vertebrate phosphodiesterase (117-119). However, no genes encoding the two activities were isolated.

A genome sequencing project for *Chlamydomonas* is underway and a simple query of the annotated gene models at <http://genome.jgi-psf.org/cgi-bin/searchGM2.cgi?db=chlre1> with “cyclase” yields 92 hits annotated as guanylyl cyclase. I analysed the bottom twelve of this list for sequence similarity with adenylyl- or guanylyl cyclases domains. Three genes harboured all the essential residues for metal binding and catalysis and these genes were all putative soluble guanylyl cyclases. The sequence of the catalytic domain of one of the genes, genie777.3, is presented in figure 1 and the domain architecture of all three is outlined in figure 2. ESTs encoding guanylyl cyclases and cGMP phosphodiesterases were also recognized in other chlorophyte algae (120). Although there is no information on the role of cGMP signaling in algae, the projected large number of guanylyl cyclases in these organisms suggests that it could be a major one.

7. ALVEOLATES

For three branches of the eukaryote tree, the Cercozoa, Heterokonts and Excavates (figure 3) there is no information on cGMP or cAMP signaling. The remaining two, the Alveolates and Discicristates contain major human pathogens and have received much closer scrutiny. The

free-living ciliates and parasitic apicomplexans are both members of the alveolates. While the ciliates are a group of fairly innocuous fresh-water protists, the apicomplexans are a family of obligatory parasites, with the malaria parasite *Plasmodium falciparum* as its most (in)famous member. Other unsavory relatives are *Toxoplasma gondii*, a major cause of congenital birth defects and *Eimeria tenella*, the causative agent of the wide-spread poultry disease, coccidiosis.

In ciliates like *Paramecium* and *Tetrahymena*, both adenylyl- and guanylyl cyclase are regulated by ion currents across the plasmamembrane. In case of adenylyl cyclase, activation is caused by an outward K^+ current and purified adenylyl cyclase protein exhibits both catalytic activity and cation conductance (121). The *Paramecium* adenylyl cyclase gene harbours both a voltage-gated cation channel with six transmembrane helices and a cytosolic adenylyl cyclase domain. There are multiple isoforms of the gene in *Paramecium* and *Tetrahymena* (79). Less well conserved orthologues are also present in several *Plasmodium* species, *Toxoplasma gondii* and *Cryptosporidium parvum* (78, 79). The *P. falciparum* genome harbours a second putative adenylyl cyclase, PfAC2, which encodes a soluble protein with a single cyclase domain (78). The cyclase domains of all alveolate adenylyl cyclases are more similar to *Dictyostelium* sGC and mammalian sAC than to the dodecahedral adenylyl cyclases (59, 80).

The *Paramecium* and *Tetrahymena* guanylyl cyclases are Ca^{2+} /calmodulin activated enzymes, that are regulated by voltage-gated Ca^{2+} influx (122-124). A single guanylyl cyclase gene was identified in the two ciliates and two orthologous genes were found in *P. falciparum*. All genes encode large proteins with in total 22 transmembrane helices. The N-terminal set of 10 helices conforms to the P-type cation transporting ATPases (125-127). However, several permutations of essential residues make it doubtful whether the ciliate ATP-ase domain can carry out active ion transport (125). The C-terminal region shows the characteristic topology of the mammalian adenylyl cyclases with two sets of six transmembrane helices interspersed and followed by a catalytic domain. However, as is the case for *Dictyostelium* GCA, the order of the C2 and C1 cyclase domains is reversed. The identity of the enzymes as a guanylyl cyclases was borne out by heterologous expression of the holoenzyme and the catalytic domains (125, 126, 128). In ciliates, both cAMP and cGMP have distinct roles in controlling cell motility (129), while in *P. falciparum*, cGMP has been implicated in exflagellation. This crucial step in the differentiation of male gametocytes is induced by xanthurenic acid (130-132). Recent work shows that xanthurenic acid increases the activity of membrane associated guanylyl cyclase, which provides an important clue for the function of the *Plasmodium* guanylyl cyclases (133). Despite its structural similarity to the G-protein regulated mammalian adenylyl cyclases, the *Paramecium* guanylyl cyclase was not activated by forskolin or by GTP γ S (128). The *P. falciparum* guanylyl cyclases cannot be regulated by heterotrimeric G-proteins *in vivo*, because heterotrimeric G-proteins are absent from

the *P. falciparum* genome (80).

8. DISCICRISTATES

The common feature of the discicristates (figure 3) are their discoid shaped mitochondrial cristae, but other aspects of their morphology are entirely different. There are amoeboid forms, of which some, the acrasids, show a primitive form of multicellularity. The euglenoids are photosynthetic freshwater protists which use a single flagellum for movement, and the trypanosomatids are obligate parasites, that retained the flagellum and the plastid, but lost the photosynthetic enzymes of the euglenoids. cAMP signaling seems to be more prevalent in the whole group than cGMP signaling. Two homologous light-activated adenylyl cyclases were isolated from the paraflagellar body of the photosynthetic flagellate *Euglena gracilis*. These remarkable enzymes are soluble proteins with two flavin binding domains that are interspersed with two cyclase catalytic domains (figure 2). Cyclase activity is 80-fold stimulated by blue light that is captured by the flavin moiety. The subsequent production of cAMP mediates the photoavoidance response of this organism (134, 135). The adenylyl cyclases that are present in the trypanosomatids *T. cruzi*, *T. brucei* and *Leishmania donovani* do not resemble the *Euglena* enzyme. They all share the same overall structure with a large extracellular domain, a single transmembrane helix and a single intracellular cyclase domain (136-139). Although their membrane topology resembles that of the receptor guanylyl cyclases, no ligand binding and kinase homology domains are present. In this aspect they resemble *Dictyostelium* ACG. All trypanosomatid cyclases are part of multigene families, but their number is particularly large in *T. brucei*. Here, at least 6 genes are part of polycistronic units that also encode the variant surface glycoproteins (VSG), which the parasite uses to escape the host immune system. These ESAG (Expression Site Associated Genes) adenylyl cyclases are only expressed when the corresponding VSG expression site is activated. In addition, several multi-membered gene families of cyclases called GRESAG (Genes Related to ESAGs) are constitutively expressed (136, 140). Only a few of the parasite genes have been functionally characterized as adenylyl cyclases and there may well be guanylyl cyclases among the remaining multitude.

Soluble NO synthase and guanylyl cyclase activities were detected in *T. cruzi* epimastigotes. The NO donor, nitroprusside stimulates the guanylyl cyclase activity 2.5-fold (141), and both nitroprusside and 8Br-cGMP stimulate epimastigote motility (142). Although the NO-induced stimulation of the parasite guanylyl cyclase is insignificant compared to the 200-fold stimulation of mammalian solGC by NO (29, 30), the data suggest that cGMP is present in trypanosomes. In *T. brucei*, this is substantiated by the presence of a highly unusual cGMP binding protein. The TbRSU gene encodes a protein, that has all the properties of PKA-R and also interacts with a PKA-type catalytic activity. However, both cNMP binding sites of TbRSU specifically bind cGMP with a $K_D \sim 10 \mu\text{M}$, while cAMP up to 100-fold higher concentrations does

not bind at all. In addition, only cGMP can activate the associated PKA activity (143). In view of these data, there are good reasons to believe that guanylyl cyclases will also be present and their sequences may be revealed once the ongoing sequencing projects of the *T. brucei* and *T. cruzi* genomes have reached completion.

9. PROKARYOTES

The prokaryotes outrank all other forms of life in the large number and diversity of nucleotidyl cyclases. There are 5 classes of adenylyl cyclases (I, II, IV, V, VI) with structurally unique catalytic domains that are absent from eukaryotes. In addition, they display an unequalled variety of the universal class III adenylyl cyclases. A recent survey of 129 bacterial genomes identified 193 class III cyclase genes in only 29 species (144). These enzymes share conserved catalytic domains with each other and with all eukaryote adenylyl- and guanylyl cyclases. However, the bacterial cyclases additionally show an unrivalled range of combinations with other functional domains and a variety of membrane topologies. The majority has a single cyclase catalytic domain, but there are also enzymes with the characteristic C1 and C2 catalytic domains of the mammalian dodecahedral adenylyl cyclases (144). A selection of cyclases with similar domain architectures as the eukaryote cyclases are presented in figure 3.

The abundance of cAMP-producing enzymes forms a stark contrast with the presence of only a single guanylyl cyclase. This enzyme, Cya2, from the cyanobacterium *Synechocystis* harbours a single cyclase domain, 4 putative transmembrane domains and a periplasmic region with a CHASE domain. This domain is also found in *Dictyostelium* ACG, in other bacterial adenylyl cyclases and in sensor histidine kinases and is hypothesized to detect small molecules (83). Targeted inactivation of Cya2 causes a 60% reduction of cGMP in the bacterium without reducing cAMP (145). However, a direct demonstration of cGMP production by the enzyme has not yet been presented.

10. SUMMARY AND PERSPECTIVE

Guanylyl- and adenylyl cyclases exist in a broad variety of structural forms. The distribution of these forms across the major groups of eukaryote organisms seems more or less random and except between closely related taxa, no patterns of inheritance are evident. In the prokaryotes, the class III cyclases show even greater structural variety and many structural forms can be recognized as prototypes for the eukaryote enzymes. For instance, the hexahelical bacterial enzymes with a single catalytic domain may, by modification of some of the helices, have given rise to the K^+ channel of the alveolate adenylyl cyclases, or by gene duplication to the dodecahedral form of the vertebrate adenylyl cyclases. The response regulator and histidine kinase domains of the Nostoc enzyme CyaC are also found in *Dictyostelium* AcrA (figure 2). The AAA-type ATP binding domain is common to the soluble eukaryote enzymes sAC and sGC, but is also

found in soluble bacterial cyclases. An extracellular CHASE domain is a common feature of *Dictyostelium* ACG and a number of bacterial cyclases.

As opposed to a scenario where all eukaryote cyclases diverged from a single ancestral eukaryote form, many of the bacterial enzymes seemed to have found their way into the eukaryotes in parallel. Some cyclases could have come with the archeobacterial ancestor of the first eukaryotes. Others could have been acquired from the ancestral bacteria and cyanobacteria that gave rise to eukaryote organelles of endosymbiotic origin, such as the mitochondria and chloroplasts. Thereafter, the emerging eukaryote groups seemed to have elaborated in different ways on some forms and have lost the others. In some organisms this loss is particularly dramatic. For example, class III guanylyl cyclases appear to be abundant in the chlorophyte algae, but have vanished from their relatives, the flowering plants. A residual function for cGMP in ion channel regulation has been retained, but when and why were the guanylyl cyclases lost? The fungi present a similar case. Phylogenetically close to both animals and dictyostelids, which share related types of adenylyl- and guanylyl cyclases, the fungi have only a single adenylyl cyclase, with little resemblance to either its the animal- and dictyostelid counterparts. When were the other forms lost and what did they regulate? The obvious common property of higher plants and fungi, that is not shared by any other group, is their sedentary life style and the complete loss of motility in all their cells including the gametes. Because cyclic nucleotide signaling is so often implicated in control of cell movement, it was possibly the loss of motility during plant and fungal evolution that led to the demise of their guanylyl- and adenylyl cyclases.

We are only at the beginning of the post-genomic era and as the number of completely sequenced genomes increases, the immense gaps in our knowledge of the evolution of cyclic nucleotide signaling will gradually be filled. Only then can we begin to understand the diversity and versatility of this most universal and most ancient system for signal transduction.

11. ACKNOWLEDGEMENT

This work is supported by BBSRC grant 94/COD16760 and Wellcome Trust University Award Grant 057137.

12. REFERENCES

1. Tesmer, J. J., R. K. Sunahara, A. G. Gilman and S. R. Sprang: Crystal structure of the catalytic domains of adenylyl cyclase in a complex with Gs α .GTP γ S. *Science* 278, 1907-1916 (1997)
2. Zhang, G., Y. Liu, A. E. Ruoho and J. H. Hurley: Structure of the adenylyl cyclase catalytic core. *Nature* 386, 247-253 (1997)
3. Yan, S. Z., Z. H. Huang, R. S. Shaw and W. J. Tang: The conserved asparagine and arginine are essential for

catalysis of mammalian adenylyl cyclase. *J Biol Chem* 272, 12342-12349 (1997)

4. Tesmer, J. J., R. K. Sunahara, R. A. Johnson, G. Gosselin, A. G. Gilman and S. R. Sprang: Two-metal-ion catalysis in adenylyl cyclase. *Science* 285, 756-760 (1999)

5. Sunahara, R. K., A. Beuve, J. J. Tesmer, S. R. Sprang, D. L. Garbers and A. G. Gilman: Exchange of substrate and inhibitor specificities between adenylyl and guanylyl cyclases. *J Biol Chem* 273, 16332-16338 (1998)

6. Tucker, C. L., J. H. Hurley, T. R. Miller and J. B. Hurley: Two amino acid substitutions convert a guanylyl cyclase, RetGC-1, into an adenylyl cyclase. *Proc Natl Acad Sci USA* 95, 5993-5997 (1998)

7. Liu, Y., A. E. Ruoho, V. D. Rao and J. H. Hurley: Catalytic mechanism of the adenylyl and guanylyl cyclases: modeling and mutational analysis. *Proc Natl Acad Sci USA* 94, 13414-13419 (1997)

8. Tang, W. J. and J. H. Hurley: Catalytic mechanism and regulation of mammalian adenylyl cyclases. *Mol Pharmacol* 54, 231-240 (1998)

9. Chinkers, M. and D. L. Garbers: The protein kinase domain of the ANP receptor is required for signaling. *Science* 245, 1392-1394 (1989)

10. Chinkers, M., D. L. Garbers, M. S. Chang, D. G. Lowe, H. M. Chin, D. V. Goeddel and S. Schulz: A membrane form of guanylate cyclase is an atrial natriuretic peptide receptor. *Nature* 338, 78-83 (1989)

11. Wilson, E. M. and M. Chinkers: Identification of sequences mediating guanylyl cyclase dimerization. *Biochemistry* 34, 4696-4701 (1995)

12. He, X., D. Chow, M. M. Martick and K. C. Garcia: Allosteric activation of a spring-loaded natriuretic peptide receptor dimer by hormone. *Science* 293, 1657-1662 (2001)

13. Labrecque, J., J. Deschenes, N. McNicoll and A. De Lean: Agonistic induction of a covalent dimer in a mutant of natriuretic peptide receptor-A documents a juxtamembrane interaction that accompanies receptor activation. *J Biol Chem* 276, 8064-8072 (2001)

14. Potter, L. R. and T. Hunter: Guanylyl cyclase-linked natriuretic peptide receptors: Structure and regulation. *J Biol Chem* 276, 6057-6060 (2001)

15. Wedel, B. and D. Garbers: The guanylyl cyclase family at Y2K. *Annu Rev Physiol* 63, 215-233 (2001)

16. Kuhn, M: Structure, regulation, and function of mammalian membrane guanylyl cyclase receptors, with a focus on guanylyl cyclase-A. *Circ Res* 93, 700-709 (2003)

17. Field, M., L. H. Graf, Jr., W. J. Laird and P. L. Smith: Heat-stable enterotoxin of *Escherichia coli*: *in vitro* effects

Guanylyl cyclase diversity

on guanylate cyclase activity, cyclic GMP concentration, and ion transport in small intestine. *Proc Natl Acad Sci USA* 75, 2800-2804 (1978)

18. Forte, L. R. and M. G. Currie: Guanylin: a peptide regulator of epithelial transport. *FASEB J* 9, 643-650 (1995)

19. Vaandrager, A. B: Structure and function of the heat-stable enterotoxin receptor/guanylyl cyclase C. *Mol Cell Biochem* 230, 73-83 (2002)

20. Fulle, H. J., R. Vassar, D. C. Foster, R. B. Yang, R. Axel and D. L. Garbers: A receptor guanylyl cyclase expressed specifically in olfactory sensory neurons. *Proc Natl Acad Sci USA* 92, 3571-3575 (1995)

21. Olshevskaya, E. V., A. N. Ermilov and A. M. Dizhoor: Dimerization of guanylyl cyclase-activating protein and a mechanism of photoreceptor guanylyl cyclase activation. *J Biol Chem* 274, 25583-25587 (1999)

22. Yu, H., E. Olshevskaya, T. Duda, K. Seno, F. Hayashi, R. K. Sharma, A. M. Dizhoor and A. Yamazaki: Activation of retinal guanylyl cyclase-1 by Ca^{2+} -binding proteins involves its dimerization. *J Biol Chem* 274, 15547-15555 (1999)

23. Olshevskaya, E. V., A. N. Ennilov and A. M. Dizhoor: Factors that affect regulation of cGMP synthesis in vertebrate photoreceptors and their genetic link to human retinal degeneration. *Mol Cell Biochem* 230, 139-147 (2002)

24. Koesling, D., C. Harteneck, P. Humbert, A. Bosserhoff, R. Frank, G. Schultz and E. Bohme: The primary structure of the larger subunit of soluble guanylyl cyclase from bovine lung. Homology between the two subunits of the enzyme. *FEBS Lett* 266, 128-132 (1990)

25. Nakane, M., S. Saheki, T. Kuno, K. Ishii and F. Murad: Molecular cloning of a cDNA coding for 70 kilodalton subunit of soluble guanylate cyclase from rat lung. *Biochem Biophys Res Commun* 157, 1139-1147 (1988)

26. Friebe, A. and D. Koesling: Regulation of nitric oxide-sensitive guanylyl cyclase. *Circ Res* 93, 96-105 (2003)

27. Wedel, B., P. Humbert, C. Harteneck, J. Foerster, J. Malkewitz, E. Bohme, G. Schultz and D. Koesling: Mutation of His-105 in the $\beta 1$ subunit yields a nitric oxide-insensitive form of soluble guanylyl cyclase. *Proc Natl Acad Sci USA* 91, 2592-2596 (1994)

28. Friebe, A., B. Wedel, C. Harteneck, J. Foerster, G. Schultz and D. Koesling: Functions of conserved cysteines of soluble guanylyl cyclase. *Biochemistry* 36, 1194-1198 (1997)

29. Arnold, W. P., C. K. Mittal, S. Katsuki and F. Murad: Nitric oxide activates guanylate cyclase and increases guanosine 3':5'-cyclic monophosphate levels in various

tissue preparations. *Proc Natl Acad Sci USA* 74, 3203-3207 (1977)

30. Zhao, Y., P. E. Brandish, D. P. Ballou and M. A. Marletta: A molecular basis for nitric oxide sensing by soluble guanylate cyclase. *Proc Natl Acad Sci USA* 96, 14753-14758 (1999)

31. Garthwaite, J., E. Southam, C. L. Boulton, E. B. Nielsen, K. Schmidt and B. Mayer: Potent and selective inhibition of nitric oxide-sensitive guanylyl cyclase by 1H-[1,2,4]oxadiazolo[4,3-a]quinoxalin-1-one. *Mol Pharmacol* 48, 184-188 (1995)

32. Zhao, Y., P. E. Brandish, M. DiValentin, J. P. Schelvis, G. T. Babcock and M. A. Marletta: Inhibition of soluble guanylate cyclase by ODQ. *Biochemistry* 39, 10848-10854 (2000)

33. Francis, S. H. and J. D. Corbin: Cyclic nucleotide-dependent protein kinases: intracellular receptors for cAMP and cGMP action. *Crit Rev Clin Lab Sci* 36, 275-328 (1999)

34. Matulef, K. and W. N. Zagotta: Cyclic nucleotide-gated ion channels. *Ann Rev Cell Dev Biol* 19, 23-44 (2003)

35. Su, Y., W. R. G. Dostmann, F. W. Herberg, K. Durick, N.-H. Xuong, L. Ten Eyck, S. S. Taylor and K. I. Varughese: Regulatory subunit of protein kinase A: structure of deletion mutant with cAMP binding domains. *Science* 269, 807-813 (1995)

36. Zoraghi, R., J. D. Corbin and S. H. Francis: Properties and functions of GAF domains in cyclic nucleotide phosphodiesterases and other proteins. *Mol Pharmacol* 65, 267-278 (2004)

37. Aravind, L. and E. V. Koonin: The HD domain defines a new superfamily of metal-dependent phosphohydrolases. *Trends Biochem Sci* 23, 469-472 (1998)

38. Soderling, S. H. and J. A. Beavo: Regulation of cAMP and cGMP signaling: new phosphodiesterases and new functions. *Curr Opin Cell Biol* 12, 174-179 (2000)

39. Francis, S. H., I. V. Turko and J. D. Corbin: Cyclic nucleotide phosphodiesterases: relating structure and function. *Prog Nucleic Acid Res Mol Biol* 65, 1-52 (2001)

40. Singh, S., D. G. Lowe, D. S. Thorpe, H. Rodriguez, W. J. Kuang, L. J. Dangott, M. Chinkers, D. V. Goeddel and D. L. Garbers: Membrane guanylate cyclase is a cell-surface receptor with homology to protein kinases. *Nature* 334, 708-712 (1988)

41. Yu, S., L. Avery, E. Baude and D. L. Garbers: Guanylyl cyclase expression in specific sensory neurons: a new family of chemosensory receptors. *Proc Natl Acad Sci USA* 94, 3384-3387 (1997)

42. L'Etoile, N. D. and C. I. Bargmann: Olfaction and odor

discrimination are mediated by the *C. elegans* guanylyl cyclase ODR-1. *Neuron* 25, 575-586 (2000)

43. Birnby, D. A., E. M. Link, J. J. Vowels, H. Tian, P. L. Colacurcio and J. H. Thomas: A transmembrane guanylyl cyclase (DAF-11) and Hsp90 (DAF-21) regulate a common set of chemosensory behaviors in *Caenorhabditis elegans*. *Genetics* 155, 85-104 (2000)

44. Morton, D. B: Invertebrates yield a plethora of atypical guanylyl cyclases. *Mol Neurobiol* 29, 97-116 (2004)

45. Morton, D. B. and A. Nighorn: MsGC-II, a receptor guanylyl cyclase isolated from the CNS of *Manduca sexta* that is inhibited by calcium. *J Neurochem* 84, 363-372 (2003)

46. Shah, S. and D. R. Hyde: Two *drosophila* genes that encode the alpha and beta subunits of the brain soluble guanylyl cyclase. *J Biol Chem* 270, 15368-15376 (1995)

47. Zayas, R. M., S. Qazi, D. B. Morton and B. A. Trimmer: Neurons involved in nitric oxide-mediated cGMP signaling in the tobacco hornworm, *Manduca sexta*. *J Comp Neurol* 419, 422-438 (2000)

48. Simpson, P. J., A. Nighorn and D. B. Morton: Identification of a novel guanylyl cyclase that is related to receptor guanylyl cyclases, but lacks extracellular and transmembrane domains. *J Biol Chem* 274, 4440-4446 (1999)

49. Nighorn, A., K. A. Byrnes and D. B. Morton: Identification and characterization of a novel beta subunit of soluble guanylyl cyclase that is active in the absence of a second subunit and is relatively insensitive to nitric oxide. *J Biol Chem* 274, 2525-2531 (1999)

50. Morton, D. B. and E. J. Anderson: MsGC-beta3 forms active homodimers and inactive heterodimers with NO-sensitive soluble guanylyl cyclase subunits. *J Exp Biol* 206, 937-947 (2003)

51. Morton, D. B., M. L. Hudson, E. Waters and M. O'Shea: Soluble guanylyl cyclases in *Caenorhabditis elegans*: NO is not the answer. *Curr Biol* 9, R546-547 (1999)

52. Bargmann, C. I: Neurobiology of the *Caenorhabditis elegans* genome. *Science* 282, 2028-2033 (1998)

53. D'Souza, C. A. and J. Heitman: Conserved cAMP signaling cascades regulate fungal development and virulence. *FEMS Microbiol Rev* 25, 349-364 (2001)

54. Borges-Walmsley, M. I. and A. R. Walmsley: cAMP signalling in pathogenic fungi: control of dimorphic switching and pathogenicity. *Trends Microbiol* 8, 133-141 (2000)

55. Casperson, G. F., N. Walker and H. R. Bourne: Isolation of the gene encoding adenylate cyclase in

Saccharomyces cerevisiae. *Proc Natl Acad Sci USA* 82, 5060-5063 (1985)

56. Toda, T., I. Uno, T. Ishikawa, S. Powers, T. Kataoka, D. Broek, S. Cameron, J. Broach, K. Matsumoto and M. Wigler: In yeast, Ras proteins are controlling elements of adenylate-cyclase. *Cell* 40, 27-36 (1985)

57. Colombo, S., P. Ma, L. Cauwenberg, J. Winderickx, M. Crauwels, A. Teunissen, D. Nauwelaers, J. H. de Winde, M. F. Gorwa, D. Colavizza and J. M. Thevelein: Involvement of distinct G-proteins, Gpa2 and Ras, in glucose- and intracellular acidification-induced cAMP signalling in the yeast *Saccharomyces cerevisiae*. *EMBO J* 17, 3326-3341 (1998)

58. Kido, M., F. Shima, T. Satoh, T. Asato, K.-I. Kariya and T. Kataoka: Critical function of the Ras-associating domain as a primary Ras-binding site for regulation of *Saccharomyces cerevisiae* adenylate cyclase. *J Biol Chem* 277, 3117-3123 (2002)

59. Linder, J. U. and J. E. Schultz: The class III adenylate cyclases: multi-purpose signalling modules. *Cellular Signalling* 15, 1081-1089 (2003)

60. Toda, T., S. Cameron, P. Sass, M. Zoller, J. D. Scott, B. McMullen, M. Hurwitz, E. G. Krebs and M. Wigler: Cloning and characterization of BCY1, a locus encoding a regulatory subunit of the cyclic AMP-dependent protein kinase in *Saccharomyces cerevisiae*. *Mol Cell Biol* 7, 1371-1377 (1987)

61. Toda, T., S. Cameron, P. Sass, M. Zoller and M. Wigler: Three different genes in *S. cerevisiae* encode the catalytic subunits of the cAMP-dependent protein kinase. *Cell* 50, 277-287 (1987)

62. Nikawa, J., P. Sass and M. Wigler: Cloning and characterization of the low-affinity cyclic AMP phosphodiesterase gene of *Saccharomyces cerevisiae*. *Mol Cell Biol* 7, 3629-3636 (1987)

63. Sass, P., J. Field, J. Nikawa, T. Toda and M. Wigler: Cloning and characterization of the high-affinity cAMP phosphodiesterase of *Saccharomyces cerevisiae*. *Proc Natl Acad Sci USA* 83, 9303-9307 (1986)

64. Chervitz, S. A., L. Aravind, G. Sherlock, C. A. Ball, E. V. Koonin, S. S. Dwight, M. A. Harris, K. Dolinski, S. Mohr, T. Smith, S. Weng, J. M. Cherry and D. Botstein: Comparison of the complete protein sets of worm and yeast: orthology and divergence. *Science* 282, 2022-2028 (1998)

65. Galagan, J. E., S. E. Calvo, K. A. Borkovich, E. U. Selker, N. D. Read, D. Jaffe, W. FitzHugh, L.-J. Ma, S. Smirnov, S. Purcell, B. Rehman, T. Elkins, R. Engels, S. Wang, C. B. Nielsen, J. Butler, M. Endrizzi, D. Qui, P. Ianakiev, D. Bell-Pedersen, M. A. Nelson, M. Werner-Washburne, C. P. Selitrennikoff, J. A. Kinsey, E. L. Braun, A. Zelter, U. Schulte, G. O. Kothe, G. Jedd, W.

- Mewes, C. Staben, E. Marcotte, D. Greenberg, A. Roy, K. Foley, J. Naylor, N. Stange-Thomann, R. Barrett, S. Gnerre, M. Kamal, M. Kamvysselis, E. Mauceli, C. Bielke, S. Rudd, D. Frishman, S. Krystofova, C. Rasmussen, R. L. Metzenberg, D. D. Perkins, S. Kroken, C. Cogoni, G. Macino, D. Catcheside, W. Li, R. J. Pratt, S. A. Osmani, C. P. C. DeSouza, L. Glass, M. J. Orbach, J. A. Berglund, R. Voelker, O. Yarden, M. Plamann, S. Seiler, J. Dunlap, A. Radford, R. Aramayo, D. O. Natvig, L. A. Alex, G. Mannhaupt, D. J. Ebbole, M. Freitag, I. Paulsen, M. S. Sachs, E. S. Lander, C. Nusbaum and B. Birren: The genome sequence of the filamentous fungus *Neurospora crassa*. *Nature* 422, 859-868 (2003)
66. Wurster, B., K. Schubiger, U. Wick and G. Gerisch: Cyclic GMP in *Dictyostelium discoideum*. Oscillations and pulses in response to folic acid and cyclic AMP signals. *FEBS Lett* 76, 141-144 (1977)
67. Ross, F. M. and P. C. Newell: Streamers: Chemotactic mutants of *Dictyostelium discoideum* with altered cyclic GMP metabolism. *J Gen Microbiol* 127, 339-350 (1981)
68. Kuwayama, H., S. Ishida and P. J. M. Van Haastert: Non-chemotactic *Dictyostelium discoideum* mutants with altered cGMP signal transduction. *J Cell Biol* 123, 1453-1462 (1993)
69. Bosgraaf, L., H. Russcher, J. L. Smith, D. Wessels, D. R. Soll and P. J. Van Haastert: A novel cGMP signalling pathway mediating myosin phosphorylation and chemotaxis in *Dictyostelium*. *EMBO J* 21, 4560-4570 (2002)
70. Pitt, G. S., N. Milona, J. Borleis, K. C. Lin, R. R. Reed and P. N. Devreotes: Structurally distinct and stage-specific adenylyl cyclase genes play different roles in *Dictyostelium* development. *Cell* 69, 305-315 (1992)
71. Roelofs, J., H. Snippe, R. G. Kleineidam and P. J. Van Haastert: Guanylate cyclase in *Dictyostelium discoideum* with the topology of mammalian adenylyl cyclase. *Biochem J* 354, 697-706 (2001)
72. Roelofs, J., M. Meima, P. Schaap and P. J. Van Haastert: The *Dictyostelium* homologue of mammalian soluble adenylyl cyclase encodes a guanylyl cyclase. *EMBO J* 20, 4341-4348 (2001)
73. Buck, J., M. L. Sinclair, L. Schapal, M. J. Cann and L. R. Levin: Cytosolic adenylyl cyclase defines a unique signaling molecule in mammals. *Proc Natl Acad Sci USA* 96, 79-84 (1999)
74. Chen, Y., M. J. Cann, T. N. Litvin, V. Iourgenko, M. L. Sinclair, L. R. Levin and J. Buck: Soluble adenylyl cyclase as an evolutionarily conserved bicarbonate sensor. *Science* 289, 625-628 (2000)
75. Saraste, M., P. R. Sibbald and A. Wittinghofer: The P-loop--a common motif in ATP- and GTP-binding proteins. *Trends Biochem Sci* 15, 430-434 (1990)
76. Confalonieri, F. and M. Duguet: A 200-amino acid ATPase module in search of a basic function. *Bioessays* 17, 639-650 (1995)
77. Soderbom, F., C. Anjard, N. Iranfar, D. Fuller and W. F. Loomis: An adenylyl cyclase that functions during late development of *Dictyostelium*. *Development* 126, 5463-5471 (1999)
78. Muhia, D. K., C. A. Swales, U. Eckstein-Ludwig, S. Saran, S. D. Polley, J. M. Kelly, P. Schaap, S. Krishna and D. A. Baker: Multiple splice variants encode a novel adenylyl cyclase of possible plastid origin expressed in the sexual stage of the malaria parasite *Plasmodium falciparum*. *J Biol Chem* 278, 22014-22022 (2003)
79. Weber, J. H., A. Vishnyakov, K. Hambach, A. Schultz, J. E. Schultz and J. U. Linder: Adenylyl cyclases from *Plasmodium*, *Paramecium* and *Tetrahymena* are novel ion channel/enzyme fusion proteins. *Cell Signal* 16, 115-125 (2004)
80. Baker, D. A. and J. M. Kelly: Purine nucleotide cyclases in the malaria parasite. *Trends Parasitol* 20, 227-232 (2004)
81. Wu, L., R. Valkema, P. J. M. Van Haastert and P. N. Devreotes: The G protein b subunit is essential for multiple responses to chemoattractants in *Dictyostelium*. *J Cell Biol* 129, 1667-1675 (1995)
82. Roelofs, J. and P. J. Van Haastert: Characterization of two unusual guanylyl cyclases from *Dictyostelium*. *J Biol Chem* 277, 9167-9174 (2002)
83. Anantharaman, V. and L. Aravind: The CHASE domain: a predicted ligand-binding module in plant cytokinin receptors and other eukaryotic and bacterial receptors. *Trends Biochem Sci* 26, 579-582 (2001)
84. Saran, S. and P. Schaap: Adenylyl cyclase G is activated by an intramolecular osmosensor. *Mol Biol Cell* 15, 1479-1486 (2004)
85. Aravind, L: An evolutionary classification of the metallo-beta-lactamase fold proteins. *In Silico Biol* 1, 69-91 (1999)
86. Lacombe, M.-L., G. J. Podgorski, J. Franke and R. H. Kessin: Molecular cloning and developmental expression of the cyclic nucleotide phosphodiesterase gene of *Dictyostelium discoideum*. *J Biol Chem* 261, 16811-16817 (1986)
87. Meima, M. E., K. E. Weening and P. Schaap: Characterization of a cAMP-stimulated cAMP phosphodiesterase in *Dictyostelium discoideum*. *J Biol Chem* 278, 14356-14362 (2003)
88. Meima, M. E., R. M. Biondi and P. Schaap:

Identification of a novel type of cGMP phosphodiesterase that is defective in the chemotactic *stmF* mutants. *Mol Biol Cell* 13, 3870-3877 (2002)

89. Bosgraaf, L., H. Russcher, H. Snippe, S. Bader, J. Wind and P. J. Van Haastert: Identification and characterization of two unusual cGMP-stimulated phosphodiesterases in *Dictyostelium*. *Mol Biol Cell* 13, 3878-3889 (2002)

90. Goldberg, J. M., L. Bosgraaf, P. J. M. Van Haastert and J. L. Smith: Identification of four candidate cGMP targets in *Dictyostelium*. *Proc Natl Acad Sci USA* 99, 6749-6754 (2002)

91. de Rooij, J., F. J. Zwartkruis, M. H. Verheijen, R. H. Cool, S. M. Nijman, A. Wittinghofer and J. L. Bos: Epac is a Rap1 guanine-nucleotide-exchange factor directly activated by cyclic AMP. *Nature* 396, 474-477 (1998)

92. Newton, R. P., L. Roef, E. Witters and H. Van Onckelen: Tansley Review No. 106 - Cyclic nucleotides in higher plants: the enduring paradox. *New Phytol* 143, 427-455 (1999)

93. The Arabidopsis Genome Initiative: Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408, 796-815 (2000)

94. Moutinho, A., P. J. Hussey, A. J. Trewavas and R. Malho: cAMP acts as a second messenger in pollen tube growth and reorientation. *Proc Natl Acad Sci USA* 98, 10481-10486 (2001)

95. Ludidi, N. and C. Gehring: Identification of a novel protein with guanylyl cyclase activity in *Arabidopsis thaliana*. *J Biol Chem* 278, 6490-6494 (2003)

96. Billington, T., M. Pharmawati and C. A. Gehring: Isolation and immunoaffinity purification of biologically active plant natriuretic peptide. *Biochem Biophys Res Comm* 235, 722-725 (1997)

97. Gehring, C. A., K. M. Khalid, T. Toop and J. A. Donald: Rat natriuretic peptide binds specifically to plant membranes and induces stomatal opening. *Biochem Biophys Res Comm* 228, 739-744 (1996)

98. Morse, M., G. Pironcheva and C. Gehring: AtPNP-A is a systemically mobile natriuretic peptide immunoanalogue with a role in *Arabidopsis thaliana* cell volume regulation. *FEBS Lett* 556, 99-103 (2004)

99. Pharmawati, M., S. N. Shabala, I. A. Newman and C. A. Gehring: Natriuretic peptides and cGMP modulate K⁺, Na⁺, and H⁺ fluxes in *Zea mays* roots. *Mol Cell Biol Res Commun* 2, 53-57 (1999)

100. Pharmawati, M., M. M. Maryani, T. Nikolakopoulos, C. A. Gehring and H. R. Irving: Cyclic GMP modulates stomatal opening induced by natriuretic peptides and immunoreactive analogues. *Plant Physiol Biochem* 39, 385-394 (2001)

101. Neill, S. J., R. Desikan and J. T. Hancock: Nitric oxide

signalling in plants. *New Phytol* 159, 11-35 (2003)

102. Pfeiffer, S., B. Janistyn, G. Jessner, H. Pichorner and R. Ebermann: Gaseous nitric-oxide stimulates guanosine-3',5'-cyclic monophosphate (cGMP) formation in spruce needles. *Phytochemistry* 36, 259-262 (1994)

103. Durner, J., D. Wendehenne and D. F. Klessig: Defense gene induction in tobacco by nitric oxide, cyclic GMP, and cyclic ADP-ribose. *Proc Natl Acad Sci USA* 95, 10328-10333 (1998)

104. Clarke, A., R. Desikan, R. D. Hurst, J. T. Hancock and S. J. Neill: NO way back: nitric oxide and programmed cell death in *Arabidopsis thaliana* suspension cultures. *Plant J* 24, 667-677 (2000)

105. Pagnussat, G. C., M. L. Lanteri and L. Lamattina: Nitric oxide and cyclic GMP are messengers in the indole acetic acid-induced adventitious rooting process. *Plant Physiol* 132, 1241-1248 (2003)

106. Schuurink, R. C., S. F. Shartzter, A. Fath and R. L. Jones: Characterization of a calmodulin-binding transporter from the plasma membrane of barley aleurone. *Proc Natl Acad Sci USA* 95, 1944-1949 (1998)

107. Kohler, C., T. Merkle and G. Neuhaus: Characterisation of a novel gene family of putative cyclic nucleotide- and calmodulin-regulated ion channels in *Arabidopsis thaliana*. *Plant J* 18, 97-104 (1999)

108. Arazi, T., R. Sunkar, B. Kaplan and H. Fromm: A tobacco plasma membrane calmodulin-binding transporter confers Ni²⁺ tolerance and Pb²⁺ hypersensitivity in transgenic plants. *Plant J* 20, 171-182 (1999)

109. Maser, P., S. Thomine, J. I. Schroeder, J. M. Ward, K. Hirschi, H. Sze, I. N. Talke, A. Amtmann, F. J. M. Maathuis, D. Sanders, J. F. Harper, J. Tchieu, M. Gribskov, M. W. Persans, D. E. Salt, S. A. Kim and M. L. Gueriot: Phylogenetic relationships within cation transporter families of *Arabidopsis*. *Plant Physiol* 126, 1646-1667 (2001)

110. Hua, B. G., R. W. Mercier, R. E. Zielinski and G. A. Berkowitz: Functional interaction of calmodulin with a plant cyclic nucleotide gated cation channel. *Plant Physiol Biochem* 41, 945-954 (2003)

111. Shabb, J. B., B. D. Buzzeo, L. Ng and J. D. Corbin: Mutating protein kinase cAMP-binding sites into cGMP-binding sites. Mechanism of cGMP selectivity. *J Biol Chem* 266, 24320-24326 (1991)

112. Clough, S. J., K. A. Fengler, I. C. Yu, B. Lippok, R. K. Smith and A. F. Bent: The *Arabidopsis* dnd1 "defense, no death" gene encodes a mutated cyclic nucleotide-gated ion channel. *Proc Natl Acad Sci USA* 97, 9323-9328 (2000)

113. Kohler, C., T. Merkle, D. Roby and G. Neuhaus: Developmentally regulated expression of a cyclic

nucleotide-gated channel from *Arabidopsis* indicates its involvement in programmed cell death. *Planta* 213, 327-332 (2001)

114. Leng, Q., R. W. Mercier, W. Z. Yao and G. A. Berkowitz: Cloning and first functional characterization of a plant cyclic nucleotide-gated cation channel. *Plant Physiol* 121, 753-761 (1999)

115. Leng, Q., R. W. Mercier, B. G. Hua, H. Fromm and G. A. Berkowitz: Electrophysiological analysis of cloned cyclic nucleotide-gated ion channels. *Plant Physiol* 128, 400-410 (2002)

116. Balague, C., B. Q. Lin, C. Alcon, G. Flottes, S. Malmstrom, C. Kohler, G. Neuhaus, G. Pelletier, F. Gaymard and D. Roby: HLM1, an essential signaling component in the hypersensitive response, is a member of the cyclic nucleotide-gated channel ion channel family. *Plant Cell* 15, 365-379 (2003)

117. Pasquale, S. M. and U. W. Goodenough: Cyclic AMP functions as a primary sexual signal in gametes of *Chlamydomonas reinhardtii*. *J Cell Biol* 105, 2279-2292 (1987)

118. Goodenough, U. W.: Cyclic AMP enhances the sexual agglutinability of *Chlamydomonas* flagella. *J Cell Biol* 109, 247-252 (1989)

119. Saito, T., L. Small and U. W. Goodenough: Activation of adenylyl cyclase in *Chlamydomonas reinhardtii* by adhesion and by heat. *J Cell Biol* 122, 137-147 (1993)

120. Becker, B., N. Feja and M. Melkonian: Analysis of expressed sequence tags (ESTs) from the scaly green flagellate *Scherffelia dubia* Pascher emend. Melkonian et Preisig. *Protist* 152, 139-147 (2001)

121. Schultz, J. E., S. Klumpp, R. Benz, W. J. C. Schurhoffgoeters and A. Schmid: Regulation of adenylyl cyclase from *Paramecium* by an intrinsic potassium conductance. *Science* 255, 600-603 (1992)

122. Schultz, J. E., T. Pohl and S. Klumpp: Voltage-gated Ca^{2+} entry into *Paramecium* linked to intraciliary increase in cyclic-GMP. *Nature* 322, 271-273 (1986)

123. Schultz, J. E. and S. Klumpp: Calcium-regulated guanylyl cyclases from *Paramecium* and *Tetrahymena*. *Method Enzymol* 195, 466-474 (1991)

124. Klumpp, S., G. Kleefeld and J. E. Schultz: Calcium calmodulin-regulated guanylate-cyclase of the excitable ciliary membrane from *Paramecium* - dissociation of calmodulin by La^{3+} - calmodulin specificity and properties of the reconstituted guanylate-cyclase. *J Biol Chem* 258, 2455-2459 (1983)

125. Linder, J. U., P. Engel, A. Reimer, T. Kruger, H. Plattner, A. Schultz and J. E. Schultz: Guanylyl cyclases with the topology of mammalian adenylyl cyclases and an

N-terminal P-type ATPase-like domain in *Paramecium*, *Tetrahymena* and *Plasmodium*. *EMBO J* 18, 4222-4232 (1999)

126. Carucci, D. J., A. A. Witney, D. K. Muhia, D. C. Warhurst, P. Schaap, M. Meima, J. L. Li, M. C. Taylor, J. M. Kelly and D. A. Baker: Guanylyl cyclase activity associated with putative bifunctional integral membrane proteins in *Plasmodium falciparum*. *J Biol Chem* 275, 22147-22156 (2000)

127. Zhang, P., C. Toyoshima, K. Yonekura, N. M. Green and D. L. Stokes: Structure of the calcium pump from sarcoplasmic reticulum at 8-A resolution. *Nature* 392, 835-839 (1998)

128. Linder, J. U., T. Hoffmann, U. Kurz and J. E. Schultz: A guanylyl cyclase from *Paramecium* with 22 transmembrane spans - Expression of the catalytic domains and formation of chimeras with the catalytic domains of mammalian adenylyl cyclases. *J Biol Chem* 275, 11235-11240 (2000)

129. Bonini, N. M. and D. L. Nelson: Differential regulation of *Paramecium* ciliary motility by cAMP and cGMP. *J Cell Biol* 106, 1615-1623 (1988)

130. Kawamoto, F., R. Alejoblanc, S. L. Fleck, Y. Kawamoto and R. E. Sinden: Possible roles of Ca^{2+} and cGMP as mediators of the exflagellation of *Plasmodium-berghei* and *Plasmodium-falciparum*. *Mol Biochem Parasitol* 42, 101-108 (1990)

131. Billker, O., V. Lindo, M. Panico, A. E. Etienne, T. Paxton, A. Dell, M. Rogers, R. E. Sinden and H. R. Morris: Identification of xanthurenic acid as the putative inducer of malaria development in the mosquito. *Nature* 392, 289-292 (1998)

132. Garcia, G. E., R. A. Wirtz, J. R. Barr, A. Woolfitt and R. Rosenberg: Xanthurenic acid induces gametogenesis in *Plasmodium*, the malaria parasite. *J Biol Chem* 273, 12003-12005 (1998)

133. Muhia, D. K., C. A. Swales, W. Deng, J. M. Kelly and D. A. Baker: The gametocyte-activating factor xanthurenic acid stimulates an increase in membrane-associated guanylyl cyclase activity in the human malaria parasite *Plasmodium falciparum*. *Mol Microbiol* 42, 553-560 (2001)

134. Iseki, M., S. Matsunaga, A. Murakami, K. Ohno, K. Shiga, K. Yoshida, M. Sugai, T. Takahashi, T. Hori and M. Watanabe: A blue-light-activated adenylyl cyclase mediates photoavoidance in *Euglena gracilis*. *Nature* 415, 1047-1051 (2002)

135. Ntefidou, M., M. Iseki, M. Watanabe, M. Lebert and D. P. Hader: Photoactivated adenylyl cyclase controls phototaxis in the flagellate *Euglena gracilis*. *Plant Physiol* 133, 1517-1521 (2003)

136. Alexandre, S., P. Paindavoine, P. Tebabi, A. Pays, S.

Halleux, M. Steinert and E. Pays: Differential expression of a family of putative adenylate/guanylate cyclase genes in *Trypanosoma brucei* 43, 279-288 (1990)

137. Taylor, M. C., D. K. Muhia, D. A. Baker, A. Mondragon, P. B. Schaap and J. M. Kelly: *Trypanosoma cruzi* adenyl cyclase is encoded by a complex multigene family. *Mol Biochem Parasitol* 104, 205-217 (1999)

138. Sanchez, M. A., D. Zeoli, E. M. Klamo, M. P. Kavanaugh and S. M. Landfear: A Family of Putative Receptor-Adenylate Cyclases from *Leishmania-Donovani*. *J Biol Chem* 270, 17551-17558 (1995)

139. D'Angelo, M. A., A. E. Montagna, S. Sanguineti, H. N. Torres and M. M. Flawia: A novel calcium-stimulated adenyl cyclase from *Trypanosoma cruzi*, which interacts with the structural flagellar protein paraflagellar rod. *J Biol Chem* 277, 35025-35034 (2002)

140. Alexandre, S., P. Paindavoine, J. Hanocq-Quertier, F. Paturiaux-Hanocq, P. Tebabi and E. Pays: Families of adenylate cyclase genes in *Trypanosoma brucei*. *Mol Biochem Parasitol* 77, 173-182 (1996)

141. Paveto, C., C. Pereira, J. Espinosa, A. E. Montagna, M. Farber, M. Esteve, M. M. Flawia and H. N. Torres: The nitric oxide transduction pathway in *Trypanosoma cruzi*. *J Biol Chem* 270, 16576-16579 (1995)

142. Pereira, C., C. Paveto, J. Espinosa, G. Alonso, M. M. Flawia and H. N. Torres: Control of *Trypanosoma cruzi* epimastigote motility through the nitric oxide pathway. *J Eukaryot Microbiol* 44, 155-156 (1997)

143. Shalaby, T., M. Liniger and T. Seebeck: The regulatory subunit of a cGMP-regulated protein kinase A of *Trypanosoma brucei*. *Eur J Biochem* 268, 6197-6206 (2001)

144. Shenoy, A. R. and S. S. Visweswariah: Class III nucleotide cyclases in bacteria and archaeobacteria: lineage-specific expansion of adenyl cyclases and a dearth of guanylyl cyclases. *FEBS Lett* 561, 11-21 (2004)

145. de Alda, J., G. Ajlani and J. Houmard: *Synechocystis* strain PCC 6803 *cya2*, a prokaryotic gene that encodes a guanylyl cyclase. *J Bacteriol* 182, 3839-3842 (2000)

146. Chenna, R., H. Sugawara, T. Koike, R. Lopez, T. J. Gibson, D. G. Higgins and J. D. Thompson: Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Res* 31, 3497-3500 (2003)

147. Schultz, J., F. Milpetz, P. Bork and C. P. Ponting: SMART, a simple modular architecture research tool: identification of signaling domains. *Proc Natl Acad Sci USA* 95, 5857-5864 (1998)

148. Bateman, A., E. Birney, R. Durbin, S. R. Eddy, K. L. Howe and E. L. L. Sonnhammer: The Pfam protein families

database. *Nucleic Acids Res* 28, 263-266 (2000)

149. Krogh, A., B. Larsson, G. von Heijne and E. L. Sonnhammer: Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* 305, 567-580 (2001)

150. Baldauf, S. L., A. J. Roger, I. Wenk-Siefert and W. F. Doolittle: A kingdom-level phylogeny of eukaryotes based on combined protein data. *Science* 290, 972-977 (2000)

151. Baldauf, S. L: The deep roots of eukaryotes. *Science* 300, 1703-1706 (2003)

Key Words: Guanylyl cyclase, cGMP phosphodiesterase, Cyclic nucleotide gated channel, PKG, Adenyl cyclase, Arthropod, Nematode, Fungi, Dictyostelium, Arabidopsis, Zea mays, Chlamydomonas, Ciliate, Plasmodium falciparum, Trypanosoma, Review

Send correspondence to: Professor Pauline Schaap, School of Life Sciences, University of Dundee, MSI/WTB complex, Dow Street, Dundee DD1 5EH, UK, Tel: 44-1382-348078, Fax: 44-1283-345386, E-mail: p.schaap@dundee.ac.uk

<http://www.bioscience.org/current/vol10.htm>