

The role of programmed-1 ribosomal frameshifting in coronavirus propagation

Ewan P. Plant¹, Jonathan D. Dinman²

¹*Division of Emerging and Transfusion Transmitted Diseases, Center for Biologics Evaluation and Research, Food and Drug Administration, Bethesda, Maryland,* ²*Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, Maryland*

TABLE OF CONTENTS

1. Abstract
2. Introduction
3. Coronavirus lifecycle
4. Programmed Frameshifting
5. What is the role of -1 PRF in the coronavirus lifecycle?
6. Perspectives
7. Acknowledgements
8. References

1. ABSTRACT

Coronaviruses have the potential to cause significant economic, agricultural and health problems. The severe acute respiratory syndrome (SARS) associated coronavirus outbreak in late 2002, early 2003 called attention to the potential damage that coronaviruses could cause in the human population. The ensuing research has enlightened many to the molecular biology of coronaviruses. A programmed -1 ribosomal frameshift is required by coronaviruses for the production of the RNA dependent RNA polymerase which in turn is essential for viral replication. The frameshifting signal encoded in the viral genome has additional features that are not essential for frameshifting. Elucidation of the differences between coronavirus frameshift signals and signals from other viruses may help our understanding of these features. Here we summarize current knowledge and add additional insight regarding the function of the programmed -1 ribosomal frameshift signal in the coronavirus lifecycle.

2. INTRODUCTION

The *Coronaviridae* family is comprised of toroviruses and at least three groups of coronaviruses which, together with the *Arteriviridae* and *Roniviridae*, belong to the order *Nidovirales* (1; <http://www.ncbi.nlm.nih.gov/ICTVdb/Ictv/index.htm>). The arteriviruses and coronaviruses can cause enteric and respiratory tract infections in mammals and birds while the roniviruses infect fish. The severity of pathogenesis varies depending on viral genotype. Outbreaks sometimes result in diarrhea and loss of livestock. After the SARS (severe acute respiratory syndrome) associated coronavirus epidemic in 2002-2003 interest and research in coronaviruses increased dramatically. Unlike the previously identified human coronaviruses (HCoV-229E and HCoV-OC43) which are more commonly associated with inconsequential respiratory infection, SARS-CoV had a mortality rate of 9.6% (2; http://www.who.int/csr/sars/country/table2004_04_21/en/index.html). The resulting resurgence in interest has added to




Figure 1. Coronavirus Lifecycle. Viral entry is mediated by a spike-receptor interaction and cathepsin L mediated membrane fusion or endocytosis (A). After endocytosis the viral genome is released (B). The first phase of translation results in the production of two polyproteins (C.) The polyproteins are processed (D) and form double membraned vesicles (DMV). The DMV is the site of both minus- and plus-strand replication (E). A second phase of translation occurs from which proteins encoded in the subgenomic RNAs are made (F). New genomic RNA is packaged by nucleocapsid protein. The structural proteins are processed in the ERGIC (G). The new viral particles exit the cell via exocytosis (H). See the main text for additional detail.

decades of work from a few groups and has advanced our knowledge of the coronavirus lifecycle considerably. This article focuses on how these viruses manipulate the host protein synthetic machinery to produce proteins essential for viral replication via a mechanism called programmed -1 ribosomal frameshifting (-1 PRF). We describe salient features and how study of the frameshifting mechanism may enhance our understanding of other parts of the viral lifecycle.

3. Coronavirus Lifecycle

Successful infection of host cells requires many steps which are common to all coronaviruses (Figure 1). Typically the coronavirus spike (S) glycoprotein and/or the haemagglutinin proteins interact with a cellular receptor to mediate entry into the host cell. Coronaviruses can also enter the cell via endocytosis. While many group 1 coronaviruses interact with an aminopeptidase, the SARS coronavirus uses the angiotensin converting enzyme 2

(ACE2) and/or the C-type lectins DC/L-SIGN as the host receptors (reviewed in 3). Cathepsin L cleaves the S protein and the viral envelope fuses with the host cell membrane. The virus disassembles, releasing the genomic RNA after which a replication-transcription complex forms on double membraned vesicles (4 and references within). New genomic and subgenomic RNA (sgRNA) is produced by the unique mechanism of discontinuous transcription during which negative-strand RNA intermediates are produced (reviewed in 5). Structural and accessory proteins are translated from the plus-strand sgRNA (Figure 2). Nucleocapsid proteins (N) package the genomic RNA and are met by envelope proteins which accumulate in the ER-to-Golgi intermediate compartment (ERGIC) for assembly. After virus particles assemble they egress from the cell via exocytosis.

Viruses by definition are dependent on the host cellular machinery for replication. However, as the viral lifecycle progresses inside the cell, the virus not only




Figure 2. SARS Coronavirus Genome Organization. (A) There are nine major open reading frames in the SARS-CoV genome. The first is divided into two overlapping parts, ORF1a and ORF1b, and comprises two thirds of the genome. In addition to the full length genomic RNA, 5' nested subgenomic RNAs are present in infected cells. The sgRNAs contain the same 5' non-coding region (bold line) as the full length genomic RNA. Two polypeptides are produced from the full length RNA (shaded boxes). They are subsequently processed into functional units, non-structural proteins nsp1 through nsp16. The structural and accessory proteins are translated from the sgRNAs (shaded boxes). See text for additional detail.

usurps certain host enzymes, but it also generates proteins that are not available from the host repertoire. For example, translation of viral proteins from the initial infectious RNA utilizes the host ribosomes while generation of new virus RNA requires a virally encoded RNA dependent RNA polymerase (RDRP). Many coronavirus proteins are translated from subgenomic RNAs (sgRNAs) rather than the genomic RNA (Figure 2). One result of this is that translation of plus-strand viral message RNA into proteins must theoretically occur in at least two phases: first the genomic RNA serves as a template for production of nonstructural proteins including the RDRP; then the RDRP uses the genomic RNA as a template for the production of sgRNAs (Figure 1). The structural and accessory proteins are produced from the sgRNAs in the next phase of translation. Thus, the second phase of translation cannot occur without the production of enzymes from the first phase. It is not known if successful infection requires the presence of more than one copy of the genomic RNA.

ORF1a/b, which is translated in the first phase, is a polyprotein that is cleaved into 16 non-structural proteins (Figure 2). The nonstructural domains in ORF1a (nsp1-11) and ORF1b (nsp12-16) are defined by proteolytic cleavage sites (reviewed in 6). The functional domains suggest that most of these proteins are involved in proteolytic cleavage and production or modification of RNA. The nonstructural proteins form replication complexes on the double membraned vesicles (DMV). During RNA replication negative strand copies of the genomic RNA are made along with negative strand sgRNAs. These in turn serve as templates for the production of positive strand genomic and

sgRNAs (reviewed in 5). The ORFs encoding structural and accessory proteins are translated from sgRNAs. The nonstructural proteins remain with the replication complex on the DMV while the structural proteins migrate for assembly into viral particles (7).

Production of proteins from ORF1a/b does not follow the usual rules of translation. Two polyproteins are produced during the translation of one disjointed open reading frame. The first polyprotein is encoded entirely within ORF1a and translation terminates at the stop codon that defines ORF1a, as is typical in normal translation. However, there are signals contained within the RNA prior to the stop codon that direct a fraction of elongating ribosomes into an alternative reading frame, allowing them to bypass the ORF1a termination codon and continue translation into ORF1b, creating a larger polyprotein. This redirection of the ribosomes to create two polyproteins has been demonstrated for many viruses including arteriviruses (8), roniviruses (9) and a number of coronaviruses (10-13). The mechanism by which the ribosomes are redirected is called programmed -1 ribosomal frameshifting (-1 PRF). It is often at least 2-orders of magnitude more efficient than baseline rates ribosomal error. The efficiency of a -1 PRF may range from 15-60% depending on the assay system and the amount of RNA flanking the core sequence (14-16). This suggests that the flanking sequences are of some importance; however, codon and reading frame constraints pose some limitations on analyses of these flanking sequences using current *in vitro* assay systems. Until the relatively recent emergence of a variety of molecular and viral tools specific to coronaviruses the pursuit of these issues has been limited. The following section describes




Figure 3. Programmed -1 Ribosomal Frameshifting. During translation features intrinsic to the mRNA being decoded manipulate the ribosome such that the reading frame is altered. (A) The ribosome pauses over a heptameric slippery site (UUUAAAC in coronaviruses). The aminoacyl-tRNA (with the filled circle) and the peptidyl-tRNA (with open circles) are positioned in the zero frame in the A- and P-sites of the ribosome. The pause and curbing of ribosome fidelity is stimulated by a 3' RNA structure. (B) Both peptidyl- and aminoacyl-tRNAs un-pair from the mRNA and re-pair in the -1 reading frame. This is facilitated by the anticodons being able to base pair at the non-wobble positions in the new reading frame. The pseudoknot structure is resolved and translation continues in the new reading frame.

how our understanding of -1 PRF has advanced with particular emphasis on the many contributions made by analysis of coronavirus frameshift signals.

4. Programmed Frameshifting

Programmed -1 ribosomal frameshifting (-1 PRF) is a mechanism used to regulate gene expression at the level of protein synthesis. As ribosomes translate one ORF they encounter a signal in the mRNA that directs a fraction of them to shift into an alternative downstream ORF which is in the -1 phase relative to the initiating upstream ORF (Figure 3). In viruses -1 PRF usually results in a C-terminally extended polypeptide containing additional function not present in the upstream ORF. The use of a -1 PRF mechanism for expression of a viral gene was first published in 1985 for the Rous sarcoma virus (17) and subsequently for other retroviruses (18). The first complete coronavirus sequence was published in 1987 (IBV; 19) and later that same an *in vitro* translation system was used to demonstrate that a -1 PRF mechanism was used to translate ORF1ab (10). In subsequent years, the IBV frameshift signal has been extensively analyzed by the Brierley and co-workers to become one of the most well characterized -1 PRF signals.

-1 PRF signals are usually composed of a “slippery site” followed by a stimulatory structure. These two elements are typically separated by a short spacer region. The slippery site is composed of a heptameric sequence such that the A- and P-site tRNAs can un-pair from the mRNA and re-pair in the -1 reading frame (20; Figure 3). The nucleotides surrounding the heptameric slippery site have been shown to have a limited effect on frameshifting efficiencies. Experiments altering the spacer region between the slippery site and stimulatory element reduced frameshifting efficiency suggesting that there

might be some optimal spacer sequence (27-30). The three nucleotides 5' of the heptameric sequence also affect -1 PRF efficiency suggesting a role for the exiting tRNA in the ribosomal E-site (24, 31). The stimulatory element has been shown to contribute significantly to -1 PRF efficiencies.

While the stimulatory structure was initially postulated to be a simple mRNA stem-loop studies of the IBV -1 PRF signal provided the first evidence for the requirement of a more complex mRNA pseudoknot (27; Figure 4). Subsequently mRNA pseudoknots were identified in the frameshift signals of a wide variety of plant and animal viruses. As additional viral sequences became available more elaborate stimulatory structures were identified in coronaviruses. These include “kissing loops” (32), and three stemmed mRNA pseudoknots, which were predicted for the coronavirus and the related torovirus Berne virus (33-34), and subsequently demonstrated by nuclease mapping for the SARS coronavirus (15-16). The variation in these stimulatory elements suggests that the additional features might be required for fine-tuning frameshifting efficiency or, alternatively, involved in additional viral functions. Interestingly, efficient frameshifting was observed when the third stem was deleted from the SARS-CoV pseudoknot, or when a similar region was deleted from the IBV stimulatory structure, suggesting that these regions are not required to modulate -1 PRF (15, 35). However, it is clear from mutational analyses that when the third stem is present that it has an effect on -1 PRF (14-15). Furthermore, additional sequence upstream of the core frameshift signal has been shown to affect -1 PRF efficiency in SARS-CoV (16). Thus, although core essential elements of the frameshift signal have been defined, the scope of factors, either cis- or trans-acting, has not yet been revealed.




Figure 4. Coronavirus Frameshift Signals. Two to three RNA stems following a heptameric slippery site comprise the coronavirus frameshift signal. The slippery site is underlined. The proposed IBV, SARS-CoV and HCoV-229E pseudoknot structures are shown (A to C respectively). The IBV structure has two stems, the SARS-CoV structure contains an additional internal stem, and the HCoV-229E structure is formed by two ‘kissing’ stem-loops. See text for additional detail.

A number of models have been proposed to describe the mechanism by which -1 PRF occurs (20-24). All the models posit that the stimulatory element causes a pause in translation and that base-pairing is required at the non-wobble positions of at least two tRNA molecules to the mRNA after the frameshift (Figure 3). Differences among the models are centered on the timing of the frameshift within the context of the elongation cycle. The detection of two different frameshift products by protein sequencing (18, 20) suggests that the different models may not be mutually exclusive. Analysis of frameshifting is complicated somewhat by the availability of malleable experimental systems imitating the appropriate host cell. It has been shown that prokaryotic ribosomes decipher coronavirus frameshift signals quite differently from yeast, plant or mammalian ribosomes (25-26). Thus a suitable system must be used to draw purposeful conclusions from *in vitro* analyses of -1 PRF. The prevalence of coronaviruses and their spread among a wide range of mammals including bats (1) suggests that analyses in mammalian cells are appropriate in most instances.

5. WHAT IS THE ROLE OF -1 PRF IN THE CORONAVIRUS LIFECYCLE?

While progress has been made on elucidating the mechanism of -1 programmed ribosomal frameshifting and the RNA sequences involved in coronaviruses, the requirement for -1 PRF in the lifecycle of this class of viruses remains obscure. For other viruses, such as HIV and the yeast totivirus L-A, frameshifting regulates the

relative ratios of structural to enzymatic proteins. The relative abundance of the coat proteins to viral RNA affects packaging and deviations from the optimal ratio result in a loss of infectivity (36-38). In contrast, -1PRF in coronaviruses modulates the relative ratios of two different classes of enzymatic proteins: proteases (and other uncharacterized proteins) encoded by the upstream Orf1a, and RDRPs and RNA modifying enzymes encoded by Orf1b. As coronavirus structural proteins are encoded on sgRNAs (the transcription of which is dependent on the frameshift), the role of -1 PRF on both the levels and timing of their synthesis, and on virus propagation in general has not yet been characterized. More specifically, the functional domains of the predicted proteases are encoded in nsp3 and nsp5 within ORF1a, prior to the -1 PRF site. The RNA modifying functions (RNA dependent RNA polymerase, helicase, exoribonuclease, uridylate-specific endoribonuclease and S-adenosylmethionine-dependent ribose 2'-O-methyltransferase) are encoded in nsp12-16 after the -1 PRF site (Figure 2). The reason for regulating the abundance of these proteins relative to one another is unknown. Nsp8 was recently described as a second RDRP raising the possibility that nsp8 and nsp12 ratios are important for controlling the amount of different RNA transcripts made during replication (39). The mechanisms by which a 100-fold more plus-strand RNA is made relative to the negative strand, or the mechanism that directs production of sgRNA rather than genomic RNA to be produced are not known (5). While it is possible that the ratio of nsp8 to nsp12 protein products may affect one of these mechanisms, this suggestion does not take into

The role of programmed-1 ribosomal frameshifting in coronavirus propagation

account the relative ratios of the other 14 proteins encoded in ORF1a/b.

The genomic RNA from plus-strand RNA viruses serves at least two functions during infection: 1) it acts as the mRNA from which viral proteins are translated, and 2) it is the template from which new genomic and subgenomic RNA is transcribed. How the infectious RNA transitions from one function to the other remains unanswered. Some progress has been made in our understanding of how another plus-strand RNA virus, the Barley yellow dwarf virus (BYDV) regulates ribosome and replicase traffic on an RNA template (40). Unlike the SARS coronavirus which utilizes a pseudoknot for frameshifting, BYDV requires a kissing-loop interaction similar to that described for the human coronavirus 229E frameshifting (32). The model requires disruption of long range RNA:RNA interactions which, if they do not reform, allow a switch between translation and transcription. In BWYV, interactions between a sgRNA and the genomic RNA also inhibit translation of the genomic RNA leaving it available for transcription or packaging (41). Such long range RNA:RNA interactions or interactions between gRNA and sgRNAs have not yet been identified in coronaviruses. An RNA switch in the 3' UTR of MHV has been characterized and found to be essential (42). This motif is found in all group 2 coronavirus sequences but only the 5' or 3' portion appears to be conserved in group 1 or group 3 coronaviruses. Interestingly a third stem-loop in the pseudoknot of the -1 PRF signal is predicted to be conserved among the group 2 coronavirus but not in the group 1 coronaviruses which utilize a kissing-loop for frameshifting (15). Some alterations to the third stem in the SARS coronavirus pseudoknot result in a loss of infectivity without dramatically affecting frameshifting, and a subset of viral proteins encoded by the subgenomic RNAs have been identified that bind to the pseudoknot in the SARS -1 PRF signal (Plant and Dinman, unpublished data). These findings suggest that this region of the SARS (+) strand is vital for an aspect of the virus lifecycle other than -1 PRF.

One current research challenge lies in producing mutations having only moderate effects on -1 PRF so that more meaningful virology can be pursued. As these mutant viruses and replicons become available we will be able to correlate the efficiency of frameshifting with production of genomic and subgenomic RNAs, and with viral titers. It is expected that some of these mutations will result in defects that will give insight into the function of the internal stem loop (stem 3) of the frameshift signal, and that that the insight thus gained will provide an alternative starting point for dissecting the coronavirus replication system.

6. PERSPECTIVES

The prospects for studies of both -1 PRF and coronaviruses are encouraging. A number of synergistic advances are being made in both areas. The constantly expanding number of coronavirus sequences is enhancing the ability of researchers to identify conserved RNA sequence and structural motifs with a greater degree of confidence. As critical elements are identified experiments

can be thoughtfully designed to generate mutant viruses and replicons from which useful information can be acquired. The size of the coronavirus genome and the lack of unique restriction sites for cloning pose some difficulties in manipulation of the virus. However recent advances in the stability of plasmid vectors and the availability of class II restriction endonucleases or restriction enzymes that cleave adjacent to the recognition sequence have circumvented some of these difficulties (43). The available clones and replicons (44-45) have allowed many groups to readily investigate different aspects of the coronavirus lifecycle. Developments in the NMR field are also enabling the solution of larger RNA structures such as those that direct -1 PRF. In addition, advances are being made in the design of algorithms able to predict tertiary RNA structures such as frameshift-promoting mRNA pseudoknots (46-48). As more structural data are generated the computational algorithms will be refined which in turn will provide enhanced tools for experimental design by bench-based researchers.

The coronavirus frameshift signals are complex and diverse and, as described above, have yielded a wealth of molecular biological data describing features important for -1 PRF. As noted previously, there are several limitations to -1 PRF studies, the most pertinent being that there are two overlapping open reading frames to maintain. Obviously silent protein coding mutations are preferable so that only recoding events are analyzed rather than protein function. The termination codon for the first ORF is very early in the SARS -1 PRF signal compared to the position of stop codons in other frameshift signals and this has increased the variety of mutations that can be sustained. A further limitation is that a frameshifting event must occur for production of the RNA dependent RNA polymerase which is essential for virus production. Thus mutations which abolish frameshifting completely will not produce replicative or infectious virus. It has been shown for some other viruses that there is an apparent threshold level of frameshifting required for competent virus production (36-38). Mutations that subtly alter the frequency of -1PRF in the coronavirus context are being discovered and these will lead to a greater understanding of the role of -1PRF in coronavirus replication.

7. ACKNOWLEDGEMENTS

Many of the published research articles would not have been possible without the free exchange of reagents and collaborations set up between groups with different expertise. The rapid emergence and severity of the SARS associated coronavirus lead to the sharing of unpublished information at conferences and meetings which in turn added vigor to the field. Because of the competitive nature of research, this sharing of resources is dependent on the ethical behavior of the researchers. We thank all those who are part of the community and have contributed important information that has the potential to prevent or control virus spread should a similar outbreak occur in the future. This work was supported by NIH grant RO1 AI064307.

The role of programmed-1 ribosomal frameshifting in coronavirus propagation

The findings and conclusions in this article have not been formally disseminated by the Food and Drug Administration and should not be construed to represent any Agency determination or policy.

8. REFERENCES

1. Vijaykrishna D, Smith GJ, Zhang JX, Peiris JS, Chen H, Guan Y: Evolutionary insights into the ecology of coronaviruses. *J Virol* 81, 4012-20 (2007)
2. Weiss SR, Navas-Martin S: Coronavirus pathogenesis and the emerging pathogen severe acute respiratory syndrome coronavirus. *Microbiol Mol Biol Rev* 69, 635-64 (2005)
3. Masters PS: The molecular biology of coronaviruses. *Adv Virus Res* 66, 193-292 (2006)
4. de Haan CA, Rottier PJ: Hosting the severe acute respiratory syndrome coronavirus: specific cell factors required for infection. *Cell Microbiol* 8, 1211-8 (2006)
5. Sawicki SG, Sawicki DL, Siddell SG: A contemporary view of coronavirus transcription. *J Virol* 81, 20-9 (2007)
6. Ziebuhr J: The coronavirus replicase. *Curr Top Microbiol Immunol* 287, 57-94 (2005)
7. Snijder EJ, van der Meer Y, Zevenhoven-Dobbe J, Onderwater JJ, van der Meulen J, Koerten HK, Mommaas AM: Ultrastructure and origin of membrane vesicles associated with the severe acute respiratory syndrome coronavirus replication complex. *J Virol* 80, 5927-40 (2006)
8. den Boon JA, Snijder EJ, Chirnside ED, de Vries AA, Horzinek MC, Spaan WJ: Equine arteritis virus is not a togavirus but belongs to the coronaviruslike superfamily. *J Virol* 65, 2910-20 (1991)
9. Cowley JA, Dimmock CM, Spann KM, Walker PJ: Gill-associated virus of *Penaeus monodon* prawns: an invertebrate virus with ORF1a and ORF1b genes related to arteri- and coronaviruses. *J Gen Virol* 81, 1473-84 (2000)
10. Brierley I, Bournsnel ME, Binns MM, Bilimoria B, Blok VC, Brown TD, Inglis SC: An efficient ribosomal frame-shifting signal in the polymerase-encoding region of the coronavirus IBV. *EMBO J* 6, 3779-85 (1987)
11. Bredenbeek PJ, Pachuk CJ, Noten AF, Charité J, Luytjes W, Weiss SR, Spaan WJ: The primary structure and expression of the second open reading frame of the polymerase gene of the coronavirus MHV-A59; a highly conserved polymerase is expressed by an efficient ribosomal frameshifting mechanism. *Nucleic Acids Res* 18, 1825-32 (1990)
12. Herold J, Raabe T, Schelle-Prinz B, Siddell SG: Nucleotide sequence of the human coronavirus 229E RNA polymerase locus. *Virology* 195, 680-91 (1993)
13. Thiel V, Ivanov KA, Putics A, Hertzog T, Schelle B, Bayer S, Weissbrich B, Snijder EJ, Rabenau H, Doerr HW, Gorbalenya AE, Ziebuhr J: Mechanisms and enzymes involved in SARS coronavirus genome expression. *J Gen Virol* 84, 2305-15 (2003)
14. Baranov PV, Henderson CM, Anderson CB, Gesteland RF, Atkins JF, Howard MT: Programmed ribosomal frameshifting in decoding the SARS-CoV genome. *Virology* 332, 498-510 (2005)
15. Plant EP, Pérez-Alvarado GC, Jacobs JL, Mukhopadhyay B, Hennig M, Dinman JD: A three-stemmed mRNA pseudoknot in the SARS coronavirus frameshift signal. *PLoS Biol* 3, e172 (2005)
16. Su MC, Chang CT, Chu CH, Tsai CH, Chang KY: An atypical RNA pseudoknot stimulator and an upstream attenuation signal for -1 ribosomal frameshifting of SARS coronavirus. *Nucleic Acids Res* 33, 4265-75 (2005)
17. Jacks T, Varmus HE: Expression of the Rous sarcoma virus pol gene by ribosomal frameshifting. *Science* 230, 1237-42 (1985)
18. Jacks T, Power MD, Masiarz FR, Luciw PA, Barr PJ, Varmus HE: Characterization of ribosomal frameshifting in HIV-1 gag-pol expression. *Nature* 331, 280-3 (1988)
19. Bournsnel ME, Brown TD, Foulds IJ, Green PF, Tomley FM, Binns MM: The complete nucleotide sequence of avian infectious bronchitis virus: analysis of the polymerase-coding region. *Adv Exp Med Biol* 218, 15-29 (1987)
20. Jacks T, Madhani HD, Masiarz FR, Varmus HE: Signals for ribosomal frameshifting in the Rous sarcoma virus gag-pol region. *Cell* 55, 447-58 (1988)
21. Weiss RB, Dunn DM, Shuh M, Atkins JF, Gesteland RF: E. coli ribosomes re-phase on retroviral frameshift signals at rates ranging from 2 to 50 percent. *New Biol* 1, 159-69 (1989)
22. Farabaugh, PJ: Programmed translational frameshifting. *Microbiol Rev* 60, 103-34 (1996)
23. Plant EP, Jacobs KL, Harger JW, Meskauskas A, Jacobs JL, Baxter JL, Petrov AN, Dinman JD: The 9-A solution: how mRNA pseudoknots promote efficient programmed -1 ribosomal frameshifting. *RNA* 9, 168-74 (2003)
24. Léger M, Dulude D, Steinberg SV, Brakier-Gingras L: The three transfer RNAs occupying the A, P and E sites on the ribosome are involved in viral programmed -1 ribosomal frameshift. *Nucleic Acids Res* 35, 5581-92 (2007)
25. Brierley I, Meredith MR, Bloys AJ, Hagervall TG: Expression of a coronavirus ribosomal frameshift signal in

The role of programmed-1 ribosomal frameshifting in coronavirus propagation

- Escherichia coli*: influence of tRNA anticodon modification on frameshifting. *J Mol Biol* 270, 360-73 (1997)
26. Plant EP, Dinman JD: Comparative study of the effects of heptameric slippery site composition on -1 frameshifting among different eukaryotic systems. *RNA* 12, 666-73 (2006)
 27. Brierley I, Digard P, Inglis SC: Characterization of an efficient coronavirus ribosomal frameshifting signal: requirement for an RNA pseudoknot. *Cell* 57, 537-47 (1989)
 28. Morikawa S, Bishop DH: Identification and analysis of the gag-pol ribosomal frameshift site of feline immunodeficiency virus. *Virology* 186, 389-97 (1992)
 29. Kollmus H, Honigman A, Panet A, Hauser H: The sequences of and distance between two cis-acting signals determine the efficiency of ribosomal frameshifting in human immunodeficiency virus type 1 and human T-cell leukemia virus type II *in vivo*. *J Virol* 68, 6087-91 (1994)
 30. Naphthine S, Liphardt J, Bloys A, Routledge S, Brierley I: The role of RNA pseudoknot stem 1 length in the promotion of efficient -1 ribosomal frameshifting. *J Mol Biol* 288, 305-20 (1999)
 31. Bekaert M, Rousset JP: An extended signal involved in eukaryotic -1 frameshifting operates through modification of the E site tRNA. *Mol Cell* 17, 61-8 (2005)
 32. Herold J, Siddell SG: An 'elaborated' pseudoknot is required for high frequency frameshifting during translation of HCV 229E polymerase mRNA. *Nucleic Acids Res* 21, 5838-42 (1993)
 33. Dos Ramos F, Carrasco M, Doyle T, Brierley I: Programmed -1 ribosomal frameshifting in the SARS coronavirus. *Biochem Soc Trans* 32, 1081-3 (2004)
 34. Snijder EJ, den Boon JA, Bredenbeek PJ, Horzinek MC, Rijnbrand R, Spaan WJ: The carboxyl-terminal part of the putative Berne virus polymerase is expressed by ribosomal frameshifting and contains sequence motifs which indicate that toro- and coronaviruses are evolutionarily related. *Nucleic Acids Res* 18, 4535-42 (1990)
 35. Brierley I, Rolley NJ, Jenner AJ, Inglis SC: Mutational analysis of the RNA pseudoknot component of a coronavirus ribosomal frameshifting signal. *J Mol Biol* 220, 889-902 (1991)
 36. Dinman JD, Wickner RB: Ribosomal frameshifting efficiency and gag/gag-pol ratio are critical for yeast M1 double-stranded RNA virus propagation. *J Virol* 66, 3669-76 (1992)
 37. Hung M, Patel P, Davis S, Green SR: Importance of ribosomal frameshifting for human immunodeficiency virus type 1 particle assembly and replication. *J Virol* 72, 4819-24 (1998)
 38. Biswas P, Jiang X, Pacchia AL, Dougherty JP, Peltz SW: The human immunodeficiency virus type 1 ribosomal frameshifting site is an invariant sequence determinant and an important target for antiviral therapy. *J Virol* 78, 2082-7 (2004)
 39. Imbert I, Guillemot JC, Bourhis JM, Bussetta C, Coutard B, Egloff MP, Ferron F, Gorbalenya AE, Canard B: A second, non-canonical RNA-dependent RNA polymerase in SARS coronavirus. *EMBO J* 25, 4933-42 (2006)
 40. Barry JK, Miller WA: A -1 ribosomal frameshift element that requires base pairing across four kilobases suggests a mechanism of regulating ribosome and replicase traffic on a viral RNA. *Proc Natl Acad Sci U S A* 99, 11133-8 (2004)
 41. Shen R, Rakotondrafara AM, Miller WA: trans regulation of cap-independent translation by a viral subgenomic RNA. *J Virol* 80, 10045-54 (2006)
 42. Goebel SJ, Miller TB, Bennett CJ, Bernard KA, Masters PS: A hypervariable region within the 3' cis-acting element of the murine coronavirus genome is nonessential for RNA synthesis but affects pathogenesis. *J Virol* 81, 1274-87 (2004)
 43. Baric RS, Sims AC: Development of mouse hepatitis virus and SARS-CoV infectious cDNA constructs. *Curr Top Microbiol Immunol* 287, 229-52 (2005)
 44. Yount B, Curtis KM, Fritz EA, Hensley LE, Jahrling PB, Prentice E, Denison MR, Geisbert TW, Baric RS: Reverse genetics with a full-length infectious cDNA of severe acute respiratory syndrome coronavirus. *Proc Natl Acad Sci U S A* 100, 12995-3000 (2003)
 45. Almazán F, Dediego ML, Galán C, Escors D, Alvarez E, Ortego J, Sola I, Zuñiga S, Alonso S, Moreno JL, Nogales A, Capiscol C, Enjuanes L: Construction of a severe acute respiratory syndrome coronavirus infectious cDNA clone and a replicon to study coronavirus RNA synthesis. *J Virol* 80, 10900-6 (2006)
 46. Reeder J, Steffen P, Giegerich R: pknotsRG: RNA pseudoknot folding including near-optimal structures and sliding windows. *Nucleic Acids Res* 35, W320-4 (2007)
 47. Dawson WK, Fujiwara K, Kawai G: Prediction of RNA pseudoknots using heuristic modeling with mapping and sequential folding. *PLoS ONE* 2, e905 (2007)
 48. Jacobs JL, Belew AT, Rakauskaitė R, Dinman JD: Identification of functional, endogenous programmed -1 ribosomal frameshift signals in the genome of *Saccharomyces cerevisiae*. *Nucleic Acids Res* 35, 165-74 (2007)

The role of programmed-1 ribosomal frameshifting in coronavirus propagation

Abbreviations: SARS: severe acute respiratory syndrome; SARS-CoV: SARS coronavirus; S: Spike protein; gRNA: genomic RNA; sgRNA: subgenomic RNA; RDRP: RNA dependent RNA polymerase; ORF: open reading frame; nsp: non-structural protein; DMV: double membraned vesicle; -1 PRF: programmed -1 ribosomal frameshifting; BYDV: Barley yellow dwarf virus.

Key Words: Coronavirus, SARS, Frameshifting, Review

Send correspondence to: Ewan Plant, CBER, FDA, HFM-310, 8800 Rockville Pike, Bethesda MD 20892, Tel: 301-827-1954, Fax: 301-480-4757, E-mail: Ewan.Plant@FDA.hhs.gov

<http://www.bioscience.org/current/vol13.htm>