

Düzeltilme/Erratum

Aralık 2017 sayımızda (18(6):571-576) yayımlanan "Bipolar bozuklukta mevsimselliği olan ve olmayan hastaların sosyodemografik ve klinik özelliklerinin karşılaştırılması" başlıklı yazının tabloları aşağıdadır.

The following tables are belong to the article entitled "Comparison of sociodemographic and clinical characteristics of bipolar patients with and without seasonal patterns" which has been published in December 2017 issue (18(6):571-576).

Table 1. Descriptive statistics of categorical clinical features between patients with and without seasonal pattern

	Seasonal pattern(n=63)		Non-seasonal pattern (n=111)		p
	n	%	n	%	
Gender					0.903
Male	21	33.3	36	32.4	
Female	42	66.7	75	67.6	
Subtype					0.500
Bipolar I disorder	61	96.8	105	94.4	
Bipolar II disorder	2	3.2	6	5.4	
First episode					0.005
Total	34	53.9	83	74.7	
First episode types in total					0.017
Manic	23	67.6	36	43.4	
Depressive	11	32.4	47	56.6	
Axis-I comorbidity (SCID-I) (current)					
Total	33	52.4	63	56.7	0.577
Alcohol, substance dependence	2	3.2	4	3.6	0.882
Panic disorder	1	1.6	8	7.2	0.108
Social anxiety disorder	13	20.6	19	17.1	0.565
Simple phobia	4	6.3	6	5.4	0.797
Obsessive compulsive disorder	10	15.8	29	26.1	0.119
Post-traumatic stress disorder	8	12.7	21	18.9	0.290
Generalized anxiety disorder	7	11.1	8	7.2	0.378
Anxiety disorder not otherwise specified	4	6.3	1	0.9	0.039
Somatoform disorders	0	0	5	4.5	0.087
Eating disorders	0	0	5	4.5	0.087
Predominant polarity					
Total	57	90.5	99	89.2	0.789
Predominant polarity types in total					0.144
Manic	34	59.6	72	72.7	
Depressive	15	26.3	21	21.2	
Mixed	8	14.0	6	6.1	
Presence of psychotic symptoms	48	76.2	83	74.7	0.835
Severe episode	35	55.5	60	54.1	0.848
Hipertymia	8	12.7	10	9.0	0.442
Distimia	8	12.7	0	0	<0.001
Presence of hospitalization	34	53.9	80	72.1	0.016
Attempted suicide	11	17.5	25	22.5	0.428
Family history of bipolar disorder	18	28.6	28	25.2	0.630
Family history of suicide	4	6.3	4	3.6	0.406
Response to lithium monotherapy	30/41	73.1	40/83	48.2	0.008
Response to anticonvulsant monotherapy	20/33	60.6	28/47	59.6	0.926
Response to antipsychotic monotherapy	10/16	62.5	17/21	80.9	0.211
Antidepressant use	35	55.5	62	55.8	0.969

The differences between those with and without seasonal pattern with respect to their categorical characteristics were compared using the Pearson's chi-square analysis and the differences with respect to numerical type of characteristics using the independent samples t-test. The statistical significance level was taken as $p < 0.05$.

Table 2. Descriptive statistics of numerical sociodemographic in patients with and without seasonal pattern

	Seasonal pattern (n=63) Mean±SD	Non-seasonal pattern (n=111) Mean±SD	p
Age	43.5±11.9	40.2±12.5	0.087
Age of onset	23.4±8.1	23.7±7.9	0.813
Years of illness	20.0±9.9	16.4±9.7	0.022
Mean episode interval (months)	13.8±10.7	12.6±6.8	0.424
Total number of episodes	11.6±8.9	9.0±5.9	0.041
Number of manic episodes	4.7±5.2	4.3±3.9	0.596
Number of hipomanic episodes	2.4±5.0	1.4±2.4	0.140
Number of mixed episodes	1.2±2.6	0.5±1.1	0.045
Number of depressive episodes	3.1±3.8	2.7±3.0	0.474
Number of hospitalization	3.3±2.1	3.1±2.5	0.574

The differences between those with and without seasonal pattern with respect to their categorical characteristics were compared using the Pearson's chi-square analysis and the differences with respect to numerical type of characteristics using the independent samples t-test. The statistical significance level was taken as $p < 0.05$.